Electronic Supplementary Information

Vibrational and Chiroptical Analysis of Blood Plasma for Hepatocellular Carcinoma Diagnostics

Ondřej Vrtělka,*^a Kateřina Králová,^a Markéta Fousková,^a Lucie Habartová,^a Petr Hříbek,^{b,c} Petr Urbánek^b and Vladimír Setnička^a

a. Department of Analytical Chemistry, University of Chemistry and Technology Prague, Technická 5, 166 28 Prague 6, Czech Republic
b. Military University Hospital Prague, Department of Medicine 1st Faculty of Medicine Charles University and Military University Hospital Prague, U Vojenské nemocnice 1200, 169 02 Prague 6, Czech Republic.

c. Department of Internal Medicine, University of Defense, Faculty of Military Health Sciences in Hradec Kralove, Třebešská 1575, 500 01 Hradec Králové, Czech Republic

*corresponding author: Ondrej.Vrtelka@vscht.cz



Fig. S1 PCA loadings generated from FTIR spectra for the PCs considered to be included in the classification model



Fig. S2 PCA explained variance of the FTIR spectra



Fig. S3 PCA loadings generated from ECD spectra for the PCs included in the classification model



Fig. S4 PCA explained variance of the ECD spectra



Fig. S5 PCA loadings generated from Raman spectra for the PCs included in the classification model



Fig. S6 PCA explained variance of the Raman spectra



Fig. S7 PCA loadings generated from ROA spectra for the PCs included in the classification model



Fig. S8 PCA explained variance of the ROA spectra