

Supplementary Information

Tracking the Cellular Uptake and Phototoxicity of Ru(II)-polypyridyl-1,8-naphthalimide Tröger's base conjugates

Sandra A. Bright,^{1,2#} MariaLuisa Erby,^{1#} Fergus E. Poynton,² Daniel Monteyne,³ David Pérez-Morga,^{3,4} Thorfinnur Gunnlaugsson,^{*2,5} D. Clive Williams,^{*1} and Robert B. P. Elmes^{*5,6,7}

¹School of Biochemistry and Immunology, Biomedical Sciences Institute, Trinity College, Dublin 2, Ireland. E-mail: clive.williams@tcd.ie; Tel: +353 1 8962596.

²School of Chemistry, Centre for Synthesis and Chemical Biology, Biomedical Sciences Institute, Trinity College Dublin, Dublin 2, Ireland. E-mail: gunnlaut@tcd.ie; Tel: +353 1 8963459.

³Laboratoire de Parasitologie Moléculaire, IBMM-DBM Université Libre de Bruxelles, Gosselies, Belgium

⁴Center for Microscopy and Molecular Imaging CMMI Université Libre de Bruxelles, Gosselies, Belgium

⁵ Synthesis and Solid State Pharmaceutical Centre (SSPC), University of Limerick, Ireland

⁶ Department of Chemistry, Maynooth University, National University of Ireland, Maynooth, Co. Kildare, Ireland. E-mail: robert.elmes@mu.ie; Tel: +353 1708 4615

⁷ Kathleen Lonsdale Institute for Human Health Research, Maynooth University, Maynooth, Co. Kildare, Ireland.

#Authors contributed equally




Figure S1: The radiation spectrum of the Hamamatsu L2570 200 W HgXe arc lamp with comparison to deuterium, halogen and xenon lamps. Reproduced from manufacturer's specification.