

Supporting information

Ruthenium and palladium bimetallic nanoparticles achieving functional parity with a rhodium cocatalyst for TiO₂-photocatalyzed ring hydrogenation of benzoic acid

*Kousuke Nakanishi,^a Sakae Araki,^a Kousuke Nomoto,^a Yuichi Onoue,^a Ryosuke Yagi,^a Hiroyuki Asakura,^b Atsuhiko Tanaka,^b Tsunehiro Tanaka^c and Hiroshi Kominami^{*b}*

^a Department of Molecular and Material Engineering, Graduate School of Science and Engineering, Kindai University, Kowakae, Higashiosaka, Osaka 577-8502, Japan

^b Department of Applied Chemistry, Faculty of Science and Engineering, Kindai University, Kowakae, Higashiosaka, Osaka 577-8502, Japan

^c Department of Molecular Engineering, Graduate School of Engineering, Kyoto University, Kyotodaigaku Katsura, Nishikyo-ku, Kyoto 615-8510, Japan

Table of Contents

Supplementary Figures

Figure S1. Fourier transforms of EXAFS spectra of Ru(X)-Pd(Y)/TiO₂ prepared by a one-step PD method, Pd foil, PdO, Ru metal powder and RuO₂ at the (a) Pd K-edge and (b) Ru K-edge. (k-range = 3-12 Å⁻¹)

Figure S2. Re-used experiments of photocatalytic hydrogenation of BA to CCA over Ru(1.0)-Pd(2.0)/TiO₂ (100 mg) in the presence of OA (1200 μmol) for 6 h.

Supplementary Figures




Figure S1. Fourier transforms of EXAFS spectra of Ru(X)-Pd(Y)/TiO₂ prepared by a one-step PD method, Pd foil, PdO, Ru metal powder and RuO₂ at the (a) Pd K-edge and (b) Ru K-edge. (k -range = 3-12 \AA^{-1})




Figure S2. Re-used experiments of photocatalytic hydrogenation of BA to CCA over Ru(1.0)-Pd(2.0)/TiO₂ (100 mg) in the presence of OA (1200 μmol) for 6 h.