

Supplemental Table 10. Postprandial Changes in Pulse Pressure at the Beginning and at the End of the study

Intervention	Baseline (0h)	Change after intervention			P (trend) ^{a,b}
		2h	4h	6h	
Baseline					
WFA (n=8)	47 (2.90)	-0.50 (3.21)	-3.75 (1.81)	-1.87 (1.80)	0.248 ^b
AI (n=9)	53 (2.96)	-2.33 (1.55)	-0.44 (1.66)	-2.44 (4.20)	0.697 ^b
RFA (n=12)	55 (3.13)	-5.25 (2.15)	-4.75 (5.14)	-1.92 (2.15)	0.132 ^a
6 weeks					
WFA (n=8)	43 (4.00)	3.87 (3.46)	1.62 (3.32)	2.37 (3.94)	0.544 ^a
AI (n=9)	51(4.60)	-0.11 (3.16)	0.44 (1.90)	-0.33(2.65)	0.834 ^a
RFA (n=9)	49 (4.68)	-1.11 (4.50)	1.22 (2.60)	2.33 (3.10)	0.364 ^b

Data expressed as mean (standard error) in mmHg

Intra-treatment comparison by General Linear Model with Bonferroni correction. P for: ^a quadratic trend; ^blinear trend . No significant results were obtained

Inter-treatment comparisons by ANCOVA Model adjusted by age and sex. No differences were observed.