

Supplemental Table 23. Changes in CRP after 6 weeks of treatment by gender

Variable	Treatment						Changes among treatments		
	WFA (n=40)		AI (n=40)		RFA (n=39)		AI vs WFA	RFA vs WFA	RFA vs AI
	Post-int	Change	Post-int	Change	Baseline	Change	Median (25 th -75 th)	Median (25 th -75 th)	Median (25 th -75 th)
<i>CRP, mg/dL</i>									
All	0.15 (0.06; 0.30)	0.000 (-0.05;0.09)	0.12 (0.06; 0.26)	-0.010 (-0.07;0.06)	0.18 (0.09;0.41)	-0.031 (-0.14;0.04)	0.010 (-0.14;0.21)	-0.050 (-0.26;0.07)	-0.025 (-0.25;0.11)
				P=0.460		P=0.230	P=0.365	0.035	P=0.202
Men (n=50)	0.19 (0.08; 0.38)	0.008 (-0.05;0.13)	0.12 (0.06; 0.31)	0.030 (-0.05;0.21)	0.13 (0.06; 0.35)	-0.020 (-0.05;0.09)	0.022 (0.00;0.08)	-0.028 (-0.04;0.00)	-0.010 (-0.18;0.00)
				P=0.071		P=0.959	P=0.730	P=0.341	P=0.478
Women (n=69)	0.12 (0.05; 0.22)	0.000 (-0.05;0.05)	0.12 (0.06; 0.23)	-0.015 (-0.09;0.01)	0.22 (0.11; 0.38)	-0.065 (-0.17;0.01)	-0.015 (-0.04;0.05)	-0.050 (-0.12;0.05)	-0.065 (-0.08;0.00)
				P=0.974		P=0.073	P=0.248	P=0.055	P=0.449

CRP, high sensitivity C reactive protein. AU, arbitrary units. Data expressed as median (25th -75th percentile). Wilcoxon and Mann Whitney tests.

Significant differences depicted in **bold** and borderline ones in *italic bold*.