

Supplemental Table 24. Postprandial Changes in C-Reactive P at the Beginning and at the End of the study

Intervention	Baseline (0h)	2h	4h	6h
<i>Baseline</i>				
WFA (n=8)	0.155 (0.09;0.19)	0.135 (0.07;0.21)*	0.140 (0.07;0.20)	0.160 (0.09;0.27)
AI (n=9)	0.130 (0.04;0.38)	0.160 (0.04;0.39)	0.170 (0.04;0.37)	0.160 (0.04;0.36)
RFA (n=12)	0.315 (0.08;0.75)	0.330 (0.08;0.53)	0.295 (0.08;0.68)*	0.150 (0.06;0.71)†
6 weeks				
WFA (n=8)	0.150 (0.09;0.53)	0.155 (0.09;0.52)	0.155 (0.09;0.56)	0.165 (0.08;0.50)
AI (n=9)	0.130 (0.07;0.33)	0.110 (0.07;0.39)	0.130 (0.07;0.44)	0.120 (0.07;0.39)
RFA (n=12)	0.140 (0.04;0.51)	0.155 (0.04;0.41)	0.150 (0.03;0.51)	0.130 (0.03;0.4)

Data expressed as median (25th-75th percentile).

Intra-treatment comparison by Wilcoxon test. * P < 0.1; †P<0.05 versus its baseline

Inter-treatment comparisons by Mann Whitney test. No differences were observed.