

Supplemental Table 27. Changes in Complement markers after 6 weeks of treatment

Variable	Treatment						Changes among treatments					
	WFA (n=36)		AI (n=34)		RFA (n=31)		AI vs WFA		RFA vs WFA		RFA vs AI	
	Post-int	Change	Post-int	Change	Post-int	Change	Mean (95%CI)	P	Mean (95%CI)	P	Mean (95%CI)	P
C1q, mg/dL (log)	1.22 ± 0.44	-0.465 (-0.75;-0.17)	1.03 ± 0.61	-0.520 (-0.81;-0.23)	0.93 ± 0.54	-0.826 (-1.1;-0.52)	-0.055 (-0.47;0.36)	0.793	-0.361 (-0.78; 0.06)	0.092 <i>con</i> <i>four</i>	-0.307 (-0.72;0.11)	0.149
C3, mg/dL (log)	1.76 ± 0.79	-0.219 (-0.64; 0.20)	1.48 ± 0.82	-0.469 (-0.89;-0.05)	1.59 ± 0.91	-0.553 (-0.99;-0.12) Igual con four	-0.251 (-0.85;0.34)	0.406	-0.335 (-0.94;0.27)	0.277	-0.084 (-0.69;0.52)	0.783
C4, mg/dL (log)	2.43 ± 0.77	-0.315 (-0.61;-0.01)	2.11 ± 0.84	-0.534 (-0.83;-0.24)	2.01 ± 0.60	-0.610 (-0.92;-0.30)	-0.220 (-0.64;0.20)	0.306	-0.296 (-0.73;0.14)	0.178	-0.076 (-0.50;0.35)	0.724 0.118 c four

Factor B, mg/dL (log)	2.43 ± 0.77	-0.352	2.11 ± 0.84	-0.451	2.01 ± 0.60	-0.565	-0.099	0.580	-0.213	0.250	-0.114	0.531
		(-0.60;-0.10)		(-0.70;-0.20)		(-0.83;-0.30)	(-0.45;0.25)		(-0.58;0.15)		(-0.47;0.25)	
											No cambio	
											con four	

Post-int, post-treatment values. Change, change from baseline. ICAM-1, intercellular adhesion molecule 1; VCAM-1, vascular cell adhesion molecule 1. Data expressed as mean ± standard deviation, mean (95% Confidence Interval, CI) or ^amedian (25th – 75th percentile). ANCOVA Model adjusted by sex and age or ^aWilcoxon and Man Whitney tests. Significant differences depicted in **bold** and borderline ones in *italic bold*.