

Supplemental Table 7b. Changes in blood pressure variables adjusted by differences in sodium consumption after 6 weeks of treatment

Variable	Treatment						Changes among treatments					
	WFA (n=37)		AI (n=38)		RFA (n=32)		AI vs WFA		RFA vs WFA		RFA vs AI	
	Post-int	Change	Post-int	Change	Post-int	Change	Mean (95%CI)	P	Mean (95%CI)	P	Mean (95%CI)	P
SBP, mmHg	121 ± 18.6 (-8.5;0.42)	-4.26	122 ± 15.2 (-4.3; 4.8)	0.305	124 ± 16.8 (-9.7; 0.30)	-4.71	4.57 (-2.0;11)	0.171	-0.448 (-7.4;6.5)	0.898	-5.02 (-12;1.7)	0.145
DBP, cm	71 ± 11.1 (-6.5;-0.62)	-3.55	73 ± 8.9 (-2.9; 2.8)	-0.009	75 ± 10.9 (-4.8;1.5)	-1.67	3.54 (-0.56;7.6)	0.090	1.88 (-2.5;6.2)	0.393	-1.66 (-5.9;2.6)	0.437
PP	50 ± 12.3 (-3.7; 2.3)	-0.718	49 ± 10.3 (-2.6; 3.2)	0.314	48 ± 11.5 (-6.3; 0.18)	-3.04	1.03 (-3.2;5.2)	0.629	-2.32 (-6.8;2.1)	0.304	-3.36 (-7.7;0.99)	0.129

Post-int, post-treatment values. Change, change from baseline. SBP, systolic blood pressure; DBP; diastolic blood pressure; PP, pulse pressure. Data expressed as mean ± standard deviation or mean (95% Confidence Interval, CI). ANCOVA Model adjusted by sex, age and differences in sodium consumption.