

Supplemental Table 7. Changes in Blood and Pulse Pressure after 6 weeks of treatment

Treatment							Changes among treatments					
Variable	WFA (n=41)		AI (n=40)		RFA (n=43)		AI vs WFA		RFA vs WFA		RFA vs AI	
	Post-int	Change	Post-int	Change	Post-int	Change	Mean (95%CI)	P	Mean (95%CI)	P	Mean (95%CI)	P
Model 1. Using V1 and V4IMP												
SBP	121 ± 18.6	-4.33 (-8.7;0.03)	122 ± 15.2	-0.021 (-4.3; 4.3)	124 ± 16.8	-4.75 (-9.2; 0.30)	4.31 (-1.8;10)	0.167	-0.416 (-6.7;5.8)	0.895	-4.72 (-11;1.4)	0.131
DBP	71 ± 11.1	-3.45 (-6.2;-0.73)	73 ± 8.9	-0.139 (-2.8; 2.5)	75 ± 10.9	-1.66 (-4.4;1.1)	3.32 (-0.52;7.1)	0.090	1.79 (-2.1;5.7)	0.364	-1.52 (-5.4;2.3)	0.435
PP	50 ± 12.3	-0.858 (-3.7; 1.9)	49 ± 10.3	0.151 (-2.6; 2.9)	48 ± 11.5	-2.95 (-5.8; -0.05)	1.01 (-2.9;4.9)	0.613	-2.09 (-6.1;2.0)	0.309	-3.10 (-7.1;0.88)	0.126

Post-int, post-treatment values. Change, change from baseline. SBP, systolic blood pressure; DBP; diastolic blood pressure; PP, pulse pressure. Data expressed as mean ± standard deviation or mean (95% Confidence Interval, CI. ANCOVA Model adjusted by sex and age. Significant differences depicted in **bold** and borderline ones in *italic bold*.