

Supplementary material

Figure S1. Dosing methods of different groups.




Table S1. Primer sequences used for RT-qPCR

Gene	Direction	Primer sequences
<i>Actin-1</i>	Forward	TCTGGATCGGAGGATCTATTCT
	Reverse	AAGCACTTTCGGTGAACAATC
<i>daf-2</i>	Forward	TGAAAGCGAAGCAGCGAGAAGG
	Reverse	CGTCCGAACTTCCGCATCACTC
<i>daf-16</i>	Forward	AATTCCTCTCAACAGCAGCAGACC
	Reverse	ACCACCGCCTTGTGACAGATTAAG
<i>skn-1</i>	Forward	GGTCTCCGTTGGCGTGATGATC
	Reverse	CTGGTGGATGCTCGGTGAGTATTG
<i>gst-24</i>	Forward	GCTCCCATTCCTCTCAGTCG
	Reverse	TGATGAGTTGGCGTAGTGGG
<i>hsp-70</i>	Forward	AGCCCGTTGTTGAGGTTGAAGTG
	Reverse	CAGGCACAGTAATGACAGCATCCC
<i>hsp-16.2</i>	Forward	CTGCAGAATCTCTCCATCTGAGTC
	Reverse	AGATTCTGAAGCAACTGCACC
<i>hsp-16.1</i>	Forward	GGCTCAGATGGAACGTCAA
	Reverse	TGGCAAACTTTTGATCATTGTTA
<i>hsp-16.11</i>	Forward	TGTTGATGTTGGTGCAGTTGCT

	Reverse	GGGCGCTTGCTGAATTGGAA
<i>age-1</i>	Forward	TCCATCTGTGGGCTCCTGAACC
	Reverse	CGCTGCGTTGGTGCCTATCC
<i>fat-5</i>	Forward	GCCCTCTTCCGTTACTGCTTCAC
	Reverse	CTTCTCCGACTGCCGCAATAGATG
<i>fat-6</i>	Forward	TCGGAGAGGGAGGTCACAACCTC
	Reverse	CGGTCGTAGACAAGTCCAAGAGC
