

A single-stage anion exchange separation method for Cd isotopic analysis in geological and environmental samples by MC-ICP-MS

Qiao-Hui Zhong^{1, 2, 3}, Lu Yin⁴, Jie Li^{1, 2,*}, Yue-Xing Feng^{5,7}, Neng-Ping Shen⁶, Bing-Yu Peng^{1,2,3}, Zhao-Yang Wang^{1,2,3}

- (1) State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, 510640, China;
- (2) CAS Center for Excellence in Deep Earth Science, Guangzhou, 510640, China;
- (3) College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing, 100101, China;
- (4) Hebei Key Laboratory of Strategic Critical Mineral Resources, Hebei GEO University, Shijiazhuang, 050031, China;
- (5) Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, 519082, PR China;
- (6) State Key Laboratory of Ore Deposit Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang, 550081, China;
- (7) Radiogenic Isotope Facility, School of Environment, The University of Queensland, Qld 4072, Australia

* Corresponding author: Jie Li, State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

E-mail address: jieli@gig.ac.cn (J. Li)

Fig. S1. Elution curves of Cd purification procedures using different reagents. Corresponding to Fig. 1a-1d, only Cd and its recovery rate at each step is included.

