


## Concomitant ion matrix effects in SCGD-OES enhanced with formic acid.

Yinchenxi Zhang<sup>a</sup>, Jaime Orejas<sup>a\*</sup>, Jorge Pisonero<sup>a\*</sup> and Nerea Bordel<sup>a</sup>

<sup>a</sup>: Grupo de Espectroscopía, Láseres y Plasmas (GELP), Department of Physics, University of Oviedo, C/ Gonzalo Gutierrez Quiros S/N, Mieres 33600, Spain

Corresponding author: [orejasjaime@uniovi.es](mailto:orejasjaime@uniovi.es); [pisonerojorge@uniovi.es](mailto:pisonerojorge@uniovi.es)



**Fig.S1** HCOOH influence on CVG-prone elemental emission signal spatially resolved pattern, from 0.5 mm above the tungsten anode to 0.5 mm below the solution surface

Table S1: The reduction potential for dissolved CVG-prone elements

Reaction	Reduction Potential (V)
$\text{Li}^+ + \text{e}^- \rightarrow \text{Li}$	-3.04 V
$\text{Sr}^{2+} + 2\text{e}^- \rightarrow \text{Sr}$	-2.90 V
$\text{Na}^+ + \text{e}^- \rightarrow \text{Na}$	-2.71 V
$\text{Mg}^{2+} + 2\text{e}^- \rightarrow \text{Mg}$	-2.37 V
$\text{Zn}^{2+} + 2\text{e}^- \rightarrow \text{Zn}$	-0.76 V
$\text{Fe}^{2+} + 2\text{e}^- \rightarrow \text{Fe}$	-0.45 V
$\text{Cd}^{2+} + 2\text{e}^- \rightarrow \text{Cd}$	-0.40 V
$\text{In}^{3+} + 3\text{e}^- \rightarrow \text{In}$	-0.34 V
$\text{Pb}^{2+} + 2\text{e}^- \rightarrow \text{Pb}$	-0.13 V
$\text{Fe}^{3+} + 3\text{e}^- \rightarrow \text{Fe}$	-0.04 V
$\text{Fe}^{3+} + \text{e}^- \rightarrow \text{Fe}^{2+}$	0.77 V
$\text{Ag}^+ + \text{e}^- \rightarrow \text{Ag}$	0.80 V
$\text{Hg}^{2+} + 2 \text{e}^- \rightarrow \text{Hg}$	0.85 V