

## Support Information

### A blood-based multi-omic landscape for the molecular characterization of Kidney Stone Disease

Weibing Pan<sup>‡1</sup>, Tianwei Yun<sup>1</sup>, Xin Ouyang<sup>2</sup>, Zhijun Ruan<sup>3</sup>, Tuanjie Zhang<sup>3</sup>, Yuhao An<sup>3\*</sup>, Rui Wang<sup>3\*</sup>, Peng Zhu<sup>2\*</sup>

\*Correspondence: Yuhao An Email: anyh@szbl.ac.cn

\*Correspondence: Rui Wang Email: wangrui@szbl.ac.cn

\*Correspondence: Peng Zhu Email: gdszpsrmmyy@163.com

1. Department of Urology, Shenzhen Pingshan People's Hospital, Shenzhen, Guangdong, 518118, China.
2. Department of Laboratory Medicine, Shenzhen Pingshan People's Hospital, Shenzhen, Guangdong, 518118, China.
3. Pingshan Translational Medicine Center, Shenzhen Bay Laboratory, Shenzhen, Guangdong, 518020, China.

#### \*Corresponding authors:

Yuhao An

Shenzhen Bay Laboratory, Pingshan translational medicine center, Shenzhen, CN 518118

Email: anyh@szbl.ac.cn

Rui Wang

Shenzhen Bay Laboratory, Pingshan translational medicine center, Shenzhen, CN 518118



Email: wangrui@szbl.ac.cn



Peng Zhu

Department of Laboratory Medicine, Shenzhen Pingshan People's Hospital, Shenzhen, CN 518020

Email: gdszpsrmmyy@163.com

## Supplementary Figures





**Figure S1 A.** The proteins changed in the “MAPK signaling pathway”. **B.** The RNAs changed in the “MAPK signaling pathway”. **C.** The proteins changed in the “WNT signaling pathway”. **D.** The RNAs changed in the “WNT signaling pathway”.