

A Rapid Synthesis of Molecularly Imprinted Polymer Nanoparticles for the Extraction of Performance Enhancing Drugs (PIEDs).

Mark V. Sullivan^{ab*}, Connor Fletcher^b, Rachel Armitage^{bc}, Chester Blackburn^{ab}, Nicholas W. Turner^{ab*}

^a Department of Chemistry, Dainton Building, University of Sheffield, Brook Hill, Sheffield S3 7HF, United Kingdom.

^b Leicester School of Pharmacy, De Montfort University, The Gateway, Leicester, LE1 9BH, United Kingdom.

^c School of Archaeology and Ancient History, University of Leicester, University Road, Leicester, LE1 7RH

*n.w.turner@sheffield.ac.uk

*mark.sullivan@sheffield.ac.uk




Figure S1: FTIR spectra of the imprinted target molecules: Andarine (A), RAD-140 (B), Estradiol (C) and Gestrinone (D), obtained using a Bruker FT-IR spectrometer (alpha model).




Figure S2: Calibration plots of the target molecules: Andarine (A), RAD-140 (B), Estradiol (C) and Testosterone (D), obtained using a Nanodrop One Spectrophotometer.




Figure S3: Scatchard analysis plot for the rebinding of the target molecules to their corresponding nanoMIPS: andarine (A), RAD-140 (B), estradiol (C), testosterone (D).




Figure S4: Scatchard analysis plot for the binding of the target molecules to the control non-imprinted polymers (NIPs): andarine (A), RAD-140 (B), estradiol (C), testosterone (D).




Figure S5: Calibration plots of the target molecules: Andarine (A), RAD-140 (B), Estradiol (C) and Testosterone (D), in River Water, obtained using a Nanodrop One Spectrophotometer.