

## Supporting Information

# Synthesis, characterization and $\varepsilon$ -caprolactone polymerization property of the ferrocenyl modified half-sandwich rare-earth metal complexes

Min Li<sup>a,c</sup>, Yi Zhong<sup>c</sup>, Yunjie Luo<sup>c,\*</sup>, and Meng Deng<sup>b,\*</sup>

<sup>a</sup> Department of Basic Course, China Coast Guard Academy, Ningbo 315800, P. R.

China

<sup>b</sup> School of Materials Science and Engineering, NingboTech University, Ningbo

315100, P. R. China

<sup>c</sup> School of Materials Science and Chemical Engineering, Key Laboratory of  
Advanced Mass Spectrometry and Molecular Analysis of Zhejiang Province, Ningbo  
University, Ningbo 315211, P. R. China

## Table of Contents

**Figure S1.**  $^1\text{H}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Sc}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**1**)

**Figure S2.**  $^{13}\text{C}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Sc}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**1**)

**Figure S3.** FT-IR spectrum (KBr) of  $(\text{FcC}_5\text{Me}_4)\text{Sc}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**1**)

**Figure S4.**  $^1\text{H}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Y}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**2**)


**Figure S5.**  $^{13}\text{C}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Y}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**2**)

**Figure S6.** FT-IR spectrum (KBr) of  $(\text{FcC}_5\text{Me}_4)\text{Y}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**2**)


**Figure S7.**  $^1\text{H}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Lu}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**3**)

**Figure S8.**  $^{13}\text{C}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Lu}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**3**)


**Figure S9.** FT-IR spectrum (KBr) of  $(\text{FcC}_5\text{Me}_4)\text{Y}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**3**)




**Figure S1.**  $^1\text{H}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Sc}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**1**) in  $\text{C}_6\text{D}_6$  (298K, 500 MHz)




**Figure S2.**  $^{13}\text{C}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Sc}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**1**) in  $\text{C}_6\text{D}_6$  (298K, 125 MHz)




**Figure S3.** FT-IR spectrum (KBr) of  $(\text{FcC}_5\text{Me}_4)\text{Sc}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**1**)




**Figure S4.** <sup>1</sup>H NMR spectrum of (FcC<sub>5</sub>Me<sub>4</sub>)Y(CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>NMe<sub>2</sub>-o)<sub>2</sub> (**2**) in C<sub>6</sub>D<sub>6</sub> (298K, 500 MHz)




**Figure S5.** <sup>13</sup>C NMR spectrum of (FcC<sub>5</sub>Me<sub>4</sub>)Y(CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>NMe<sub>2</sub>-o)<sub>2</sub> (**2**) in C<sub>6</sub>D<sub>6</sub> (298K, 125 MHz)




**Figure S6.** FT-IR spectrum (KBr) of  $(\text{FcC}_5\text{Me}_4)\text{Y}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**2**)



**Figure S7.**  $^1\text{H}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Lu}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**3**) in  $\text{C}_6\text{D}_6$  (298K, 500 MHz)



**Figure S8.**  $^{13}\text{C}$  NMR spectrum of  $(\text{FcC}_5\text{Me}_4)\text{Lu}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**3**) in  $\text{C}_6\text{D}_6$  (298K, 125 MHz)



**Figure S9.** FT-IR spectrum (KBr) of  $(\text{FcC}_5\text{Me}_4)\text{Y}(\text{CH}_2\text{C}_6\text{H}_4\text{NMe}_2\text{-o})_2$  (**3**)