

L-Proline catalyzed synthesis of 2-aryl-2*H*,5*H*-thiopyrano[2,3-*b*]thiochromen-5-ones from 4-hydroxydithiocoumarins and cinnamaldehyde derivatives

Electronic Supporting Information

Ahmad Ali, Simra Faraz, and Abu Taleb Khan*

Department of Chemistry, Indian Institute of Technology Guwahati, Guwahati-781039, Assam, India.

E-mail: atk@iitg.ac.in

Table of Contents:

1. General information and methods.....	S1
2. X-ray crystallographic data of compound 3e	S2
3. Substrate scope.....	S3
4. Unsuccessful results.....	S4
5. ^1H , ^{13}C spectrum and HRMS of all compounds.....	S5-S83
6. HRMS of intermediate A1	S84

1. General information and methods

Starting material 4-Hydroxydithiocoumarin and 4-hydroxythiocoumarin was synthesize in our laboratory via previously reported methods. While cinnamaldehyde derivatives were all Sigma Aldrich. Melting points were recorded using the melting point apparatus (Stuart company). ^1H and ^{13}C NMR spectra were recorded on 400 and 500 MHz and 100, 125 MHz (Bruker) NMR spectrometers, respectively. TMS is used as an internal reference; chemical shifts (δ scale) are reported in parts per million (ppm). ^1H NMR spectra are reported in the order of multiplicity, coupling constant (J value) in Hertz (Hz), and the number of protons; signals were characterized as s (singlet), d (doublet), t (triplet), m (multiplet) and bs (broad). IR spectra were recorded on an IR spectrophotometer. HRMS spectra were recorded on a TOF mass analyser. The structure was also determined using a single XRD diffractometer.

2. X-ray crystallographic data of compound 3e





Figure. ORTEP Diagram of compound 3e

Table S1. Crystal data and structure refinement for compound (3e)

Entry	Identification Code	Compound 3e
01	Empirical formula	<u>C₁₉H₁₄O₂S₂</u>
02	Formula weight	338.42
03	Temperature	<u>297</u> K
04	Wavelength	0.71073
05	Radiation type	Mo K\alpha
06	Radiation system	Fine-focus sealed tube
07	Crystal system	<u>Triclinic</u>
08	Space group	P-1
09	Cell length(Å)	(a)7.6579 (10), (b)13.8840(19) (c)14.998 (2)
10	Cell angle (°)	α 87.216 (3), β 89.610 (4), γ 83.982 (3)
11	Cell volume (Å ³)	1584.0 (4)

12	Density	1.419
13	Absorption correction	multi-scan
14	Reflection number	6867
15	Cell formula units Z	2
16	CCDC no	2311036


3. Substrate scope^{a,b}



$R_1 = 5CL, 6CL, 7CL, 7F, 6CF3$

$R_2 = Br, -CH_3, -OMe, NO_2, -NMe_2$

$R_3 = -CH_3$




^aAll the reactions

were

carried out using 4-hydroxydithiocoumarin (**1**, 0.2 mmol), cinnamalhyde (**2**, 0.2 mmol) in the presence of 20 mol% catalyst, 1 mL of methanol Solvent at reflux temperature. ^bIsolated yield.


4. Unsuccessful results^a




The reactions were carried out using 4-hydroxydithiocoumarin (**1**, 0.2 mmol), Nitrocinnamaldehyde (**4**, 0.2 mmol) in the presence of 20 mol% catalyst, 1 mL of methanol Solvent at reflux temperature.


5. ^1H , ^{13}C spectrum and HRMS of all compounds

^1H NMR Compound (3a)







HRMS 3a




^1H NMR Compound (3b)


ATK-AA-²²FH₃
ATK-AA-²²DH₃




-3.78




¹³C NMR Compound (3b)




HRMS of compound 3b




¹H NMR Compound (3c)



ATK-AA-I2-P2A12-13C.1.fid
ATK-AA-I2-P2A12-13C




HRMS compound (3c)




MPO-AA-31-1H.5.fid
MPO-AA-31-1H


¹H NMR of compound (3d)



(500 MHz, DMSO-*d*₆) δ 8.78 (s, 1H), 7.72 (d, *J* = 8.3 Hz, 1H), 7.52 (d, *J* = 8.4 Hz, 1H), 7.29 – 7.26 (m, 3H), 6.85 (d, *J* = 8.6 Hz, 2H), 5.99 (dd, *J* = 10.2, 5.5 Hz, 1H), 4.94 (d, *J* = 5.1 Hz, 1H), 3.77 (s, 3H).






Hrms 3d




¹H NMR Compound (3e)

ATK-AA-3-P2-A3-1H.18^{fid}
ATK-AA-3-P2-A3-1H 8.51 8.51




¹³C NMR Compound (3e)

ATK-AA-3-P2-A3-13C.3.fid
ATK-AA-3-P2-A3-13C






HRMS (3e)




¹H NMR Compound (3f)

ATK-AA-23-1H.5.fid
ATK-AA-23-1H








HRMS 3f




¹H NMR Compound (3g)

ATK-AA-11-P11-1H.1.fid — ATK-AA-11-P11-1H




¹³C NMR Compound (3g)


ATK-AA-11-P11-13C.3.fid
ATK-AA-11-P11-13C




HRMS compound (3g)




¹H NMR of compound (3h)






HRMS of compound (3h)




¹H NMR of compound (3i)

ATK-AA-127-1H.1.fid
ATK-AA-127-1H




(400 MHz, Chloroform-*d*) δ 8.81 – 8.79 (m, 1H), 7.75 – 7.70 (m, 1H), 7.52 (d, *J* = 8.4 Hz, 1H), 7.38 (d, *J* = 10.3 Hz, 1H), 7.32 – 7.27 (m, 2H), 6.94 – 6.87 (m, 2H), 5.95 (dd, *J* = 10.3, 6.3 Hz, 1H), 5.35 (d, *J* = 6.2 Hz, 1H), 3.91 (s, 3H).



ATK-AA-27-13C.1.fid
ATK-AA-27-13C



HRMS 3i




¹H NMR Compound (3j)




¹³C NMR Compound (a3j)

ATK-AA-8-13C.1.fid
ATK-AA-8-13C




HRMS of 3j




¹H NMR Compound (3k)

ATK-AA-13-P2A13¹H1.fid
ATK-AA-13-P2A13¹H1.fid





¹H NMR Compound (3k)

ATK-AA-13-P2A13-13C.3.fid
ATK-AA-13-P2A13-13C




HRMS Compound (3k)








ATK-AA-24-13C.5.fid
ATK-AA-24-13C




HRMS (3l)




¹H NMR of compound (3m)







HRMS of 3m




¹H NMR Compound (n)

ATK-AA-18-1H.1.fid
ATK-AA-18-1H




¹³C NMR Compound (3n)


ATK-AA-18-13C.1.fid
ATK-AA-18-13C




HRMS of 3n




¹H NMR Compound (3o)









HRMS 3o



¹H NMR Compound (3p)





ATK-AA-14-13C.21.fid
ATK-AA-14-13C




¹³C NMR (126 MHz, Chloroform-*d*) δ 175.1, 145.3, 138.6, 136.1, 132.2 (2C), 131.5, 131.3, 129.6, 129.6 (2C), 127.8, 125.8, 125.4, 124.5, 122.6, 120.7, 43.3.

HRMS 3p







ATK-AA-19-4BR-13C.1.fid
ATK-AA-19-4BR-13C




HRMS 3q




¹H NMR of compound (3r)







HRMS of 3r



¹H NMR of compound (3s)



ATK-AA-29-1H.
ATK-AA-29-1H

ATK-AA-29-1H




¹³C NMR of compound (3s)

ATK-AA-29-13C.5.fid
ATK-AA-29-13C




HRMS a(3s)




ATK-AA-35-1H.1.fid
ATK-AA-35-1H


¹H NMR Compound (3t)



ATK-AA-35-13C.3.fid
ATK-AA-35-13C




HRMS (3t)



¹H NMR (**3u**)

ATK-AA-28-1H.1^{fid}
ATK-AA-28-1H



¹³C NMR of compound (3u)

ATK-AA-28-13C.5.fid
ATK-AA-28-13C

— 174.48

— 141.33

— 139.12

✓ 137.98

✓ 137.73



✓ 141.33
✓ 139.12
✓ 137.98
✓ 137.73
✓ 131.27
✓ 131.24
✓ 129.63
✓ 129.14
✓ 128.56
✓ 128.22
✓ 127.49
✓ 126.55
✓ 124.73
✓ 119.91
✓ 131.27
✓ 131.24
✓ 129.63
✓ 129.14
✓ 128.56
✓ 128.22
✓ 127.49
✓ 126.55

— 124.73


✓ 77.41
✓ 77.16
✓ 76.91

— 47.76

— 23.51




HRMS 3u




^1H NMR Compound (3v)


ATK-AA-15-P2P-1H.1.f1
ATK-AA-15-P2P-1H ∞





ATK-AA-15-P²⁹P-13C.3.fid
ATK-AA-15-P²⁹P-13C




HRMS (3v)




¹H NMR of compound (3w)



¹³C NMR of compound (3w)




HRMS 3w




ATK-AA20-1H.3.fid
ATK-AA20-1H


¹H NMR Compound (3x)




ATK-AA20-13C.1.fid
ATK-AA20-13C



HRMS of compound (3x)




¹H NMR of compound (3y)




¹³C NMR of compound (3y)


ATK-AA-M-CH3-13C.3.9d
ATK-AA-M-CH3-13C




HRMS of compound (3y)




¹H NMR of compound (3z)




¹³C NMR of compound (3z)



HRMS of compound (3z)



6. HRMS of intermediate (A1)



Expected m/z calculated for $\text{C}_{14}\text{H}_{16}\text{NO}_2$ 230.1176($\text{M} + \text{H}^+$); Found 230.1187.

