


Supporting Information

Development of polyethyleneimine cross-linked fucoidan nanoparticles as delivery systems for improved anticancer efficiency of cytarabine in breast adenocarcinoma cell lines.


Deepa Geethakumari ^a, Santhini Pulikkal Veettil ^{a,d}, Sivakumar Krishnankutty Nair Chandrika^{a,b}, Anoop Bhaskaran Sathyabhama ^a, Rojin Joseph ^a, Shibin Sobhanam Padmini^a, Jisha V Somasekharan ^c and Sajeevan Thavarool Puthiyedathu ^{e*}

- a. National Centre for Aquatic Animal Health, Cochin University of Science and Technology, Cochin, Kerala – 16
- b. Rajiv Gandhi Centre for Biotechnology, Poojappura, Thiruvananthapuram, Kerala - 695014
- c. Research and Post Graduate Department of Chemistry, MES Keveeyam College, Valanchery-676552, Kerala, India
- d. Department of Applied Chemistry, Cochin University of Science and Technology, Cochin, Kerala –682022, India
- e. Department of Marine Biology, Microbiology and Biochemistry, Cochin University of Science and Technology, Cochin, Kerala – 16


*Corresponding author: Sajeevan Thavarool Puthiyedathu, Tel: +91-9946099408, Fax: +91484 2381120, E-mail: sajeev@cusat.ac.in



Supplementary Fig 1. X-ray diffractograms of fucoidan (a) and fucoidan-PEI complex (b)



Supplementary Fig 2. TGA curves of fucoidan, PEI and FuNPs



Supplementary Fig 3 Cell viability studies of cytarabine and CFNP in MCF-7 cell lines by MTT assay; Results expressed as mean \pm SD (p value ≤ 0.05)