Electronic Supplementary Information

Metal-free polycyclotrimerization of trimethylsilyl-protected diynes: a facile strategy toward regioregular hyperbranched polyarylenes

Lu Wang,^{‡a} Yongliang Zhan,^{‡a} Linhui Zhang,^a and Hongkun Li,^{*a}

^aLaboratory of Advanced Optoelectronic Materials, Suzhou Key Laboratory of Novel Semiconductor-Optoelectronics Materials and Devices, State and Local Joint Engineering Laboratory for Novel Functional Polymeric Materials, Jiangsu Engineering Laboratory of Novel Functional Polymeric Materials, College of Chemistry, Chemical Engineering and Materials Science, Soochow University, Suzhou, 215123, China. E-mail: <u>hkli@suda.edu.cn</u>

‡L. Wang and Y. Zhan contributed equally to this work.

Table of Contents

Scheme S1. Synthetic route to monomer 1c

Scheme S2. Synthetic route to model compound 5

Scheme S3. Synthetic route to model compound 8

Fig. S1 ¹H NMR spectrum of intermediate compound 2 in DMSO- d_6 without TMS. The solvent and water peaks are marked with asterisks.

Fig. S2 ¹³C NMR spectra of intermediate compound 2 in DMSO- d_6 without TMS. The solvent peaks are marked with asterisks.

Fig. S3 ¹H NMR spectrum of intermediate compound **3** in CDCl₃. The solvent and water peaks are marked with asterisks.

Fig. S4 ¹³C NMR spectra of intermediate compound 3 in CDCl₃. The solvent peaks are marked with asterisks.

Fig. S5 ¹H NMR spectra of compound **4** in CDCl₃. The solvent and water peaks are marked with asterisks.

Fig. S6 ¹³C NMR spectra of compound 4 in CDCl₃. The solvent peaks are marked with asterisks. Fig. S7 IR spectra of model compound 5. **Fig. S8** ¹³C NMR spectra of model compound **5** in CD_2Cl_2 without TMS. The solvent peaks are marked with asterisks.

Fig. S9 IR spectra of model compound 8.

Fig. S10 ¹³C NMR spectra of model compound 8 in CD_2Cl_2 without TMS. The solvent peaks are marked with asterisks.

Fig. S11 IR spectra of monomer 1b (A) and its polymer *hb*-P1b (B).

Fig. S12 IR spectra of monomer 1c (A) and its polymer *hb*-P1c (B).

Fig. S13 IR spectra of monomer 1d (A) and its polymer *hb*-P1d (B).

Fig. S14 IR spectra of monomer 1e (A) and its polymer *hb*-P1e (B).

Fig. S15 ¹H NMR spectra of monomer **1b** (A) and polymer *hb*-P**1b** (B) in CD_2Cl_2 without TMS. The solvent and water peaks are marked with asterisks.

Fig. S16 ¹H NMR spectra of monomer **1c** (A) and polymer hb-P**1c** (B) in CD₂Cl₂ without TMS. The solvent and water peaks are marked with asterisks.

Fig. S17 ¹H NMR spectra of monomer **1d** (A) and polymer hb-P1d (B) in CD₂Cl₂ without TMS. The solvent and water peaks are marked with asterisks.

Fig. S18 ¹H NMR spectra of monomer **1e** (A) and polymer hb-P**1e** (B) in CD₂Cl₂ without TMS. The solvent and water peaks are marked with asterisks.

Fig. S19 ¹³C NMR spectra of monomer **1b** (A) and polymer hb-P**1b** (B) in CD₂Cl₂ without TMS. The solvent peaks are marked with asterisks.

Fig. S20 ¹³C NMR spectra of monomer **1c** (A) and polymer hb-P**1c** (B) in CD₂Cl₂ without TMS. The solvent peaks are marked with asterisks.

Fig. S21 ¹³C NMR spectra of monomer **1d** (A) and polymer hb-P**1d** (B) in CD₂Cl₂ without TMS. The solvent peaks are marked with asterisks.

Fig. S22 ¹³C NMR spectra of monomer **1e** (A) and polymer hb-P**1e** (B) in CD₂Cl₂ without TMS. The solvent peaks are marked with asterisks.



Scheme S1 Synthetic route to monomer 1c







Scheme S3 Synthetic route to model compound 8



Fig. S1 ¹H NMR spectra of intermediate compound 2 in DMSO- d_6 without TMS. The solvent and water peaks are marked with asterisks.



Fig. S2 ¹³C NMR spectra of intermediate compound 2 in DMSO- d_6 without TMS. The solvent peaks are marked with asterisks.



Fig. S3 ¹H NMR spectra of intermediate compound 3 in $CDCl_3$. The solvent and water peaks are marked with asterisks.



Fig. S4 13 C NMR spectra of intermediate compound 3 in CDCl₃. The solvent peaks are marked with asterisks.



Fig. S5 ¹H NMR spectra of compound 4 in CDCl₃. The solvent and water peaks are marked with asterisks.



Fig. S6¹³C NMR spectra of compound 4 in CDCl₃. The solvent peaks are marked with asterisks.



Fig. S7 IR spectra of model compound 5.



Fig. S8 ¹³C NMR spectra of model compound 5 in CD_2Cl_2 without TMS. The solvent peaks are marked with asterisks.



Fig. S9 IR spectra of model compound 8.



Fig. S10 ¹³C NMR spectra of model compound 8 in CD_2Cl_2 without TMS. The solvent peaks are marked with asterisks.



Fig. S11 IR spectra of monomer 1b (A) and its polymer *hb*-P1b (B).



Fig. S12 IR spectra of monomer 1c (A) and its polymer *hb*-P1c (B).



Fig. S13 IR spectra of monomer 1d (A) and its polymer hb-P1d (B).



Fig. S14 IR spectra of monomer 1e (A) and its polymer hb-P1e (B).



Fig. S15 ¹H NMR spectra of monomer **1b** (A) and polymer *hb*-P**1b** (B) in CD_2Cl_2 without TMS. The solvent and water peaks are marked with asterisks.



Fig. S16 ¹H NMR spectra of monomer **1c** (A) and polymer *hb*-P**1c** (B) in CD_2Cl_2 without TMS. The solvent and water peaks are marked with asterisks.



Fig. S17 ¹H NMR spectra of monomer **1d** (A) and polymer *hb*-P**1d** (B) in CD_2Cl_2 without TMS. The solvent and water peaks are marked with asterisks.



Fig. S18 ¹H NMR spectra of monomer **1e** (A) and polymer hb-P1e (B) in CD₂Cl₂. without TMS. The solvent and water peaks are marked with asterisks.



Fig. S19 ¹³C NMR spectra of monomer **1b** (A) and polymer *hb*-P**1b** (B) in CD_2Cl_2 without TMS. The solvent peaks are marked with asterisks.



Fig. S20 ¹³C NMR spectra of monomer **1c** (A) and polymer hb-P**1c** (B) in CD₂Cl₂ without TMS. The solvent peaks are marked with asterisks.



Fig. S21 ¹³C NMR spectra of monomer **1d** (A) and polymer hb-P1d (B) in CD₂Cl₂ without TMS. The solvent peaks are marked with asterisks.



Fig. S22 ¹³C NMR spectra of monomer **1e** (A) and polymer *hb*-P**1e** (B) in CD_2Cl_2 without TMS. The solvent peaks are marked with asterisks.