

Pathogen inhibition and indication by gelatin nonwoven mats with incorporation of polyphenol derivatives

By *Meng-Yi Bai*^{a, .b, *}, *Ting-Teng Wang*^a, *Xin-An Chen*^a, *Chia-Chun Wu*^c

[a] Graduate Institute of Biomedical Engineering and Biomedical Engineering Program, Graduate Institute of Applied Science and Technology, National Taiwan University of Science and Technology, TR-917, AAEON Building, No. 43, Keelung Rd., Sec. 4, Da'an Dist., Taipei City 10607, Taiwan, ROC

[b] Adjunct Appointment to the National Defense Medical Center, Taipei 11490, Taiwan, ROC

[c] Institute of Prevention Medicine, National Defense Medical Center, Taipei 11490, Taiwan, ROC

[*] Correspondence to: Prof. M.-Y. Bai, Ph.D.

Graduate Institute of Biomedical Engineering, National Taiwan University of Science and Technology, TR-917, AAEON Building, No. 43, Keelung Rd., Sec. 4, Da'an Dist., Taipei City 10607, Taiwan, ROC

E-mail: mybai@mail.ntust.edu.tw

This work was supported by grants from the Ministry of Science and Technology, Taiwan (MOST 109-2221-E-011-036, 108-2221-E-011-113, and 111-2221-E-011-151).




Figure S1 HPLC analysis and the chromatograms of standard compounds of (a) leucine, (b) valine, (c) glutamate, and (d) aspartate for comparison.




Figure S2 MTT assay for determining the safe concentration that was identified through a series of different concentrations (a) 125 ppm-8000 ppm, and (b) 5000-80000 rpm, against MDCK cells.