

Supporting Information for

Synthesis of *N*-glycosyl amides. Conformational analysis and evaluation as inhibitors of the β -galactosidase from *E. coli*.

Miqueas G. Traverssi,^{ab}, Verónica E. Manzano,^{cd} Oscar Varela^{cd} and Juan P. Colomer *^{ab}

^aDepartamento de Química Orgánica, Universidad Nacional de Córdoba, Facultad Ciencias Químicas, Ciudad Universitaria, Edificio de Ciencias II, Córdoba, Argentina. E-mail: juanpablo@fcq.unc.edu.ar

^bInstituto de Investigaciones en Físico-Química de Córdoba (INFIQC), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), UNC, Argentina

^cDepartamento de Química Orgánica, Universidad de Buenos Aires, Facultad Ciencias Exactas y Naturales, Ciudad Universitaria, Pab. 2, C1428EHA, Buenos Aires, Argentina

^dCentro de Investigación en Hidratos de Carbono (CIHIDECAR), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), UBA, Argentina


Index

Contents


Table 1. Chair conformations 4C_1 and 1C_4 and energies obtained for the <i>xylo</i> derivates by conformational search employing MMFF94.	2
1H and ^{13}C NMR spectra of glycosyl azides 5a-c	3
1H and ^{13}C NMR spectra of thioglyco acid 13β and 13α	9
1H and ^{13}C NMR spectra of glycosylamines 2cα and 2cβ	12
1H , ^{13}C , and NOESY NMR spectra of per- <i>O</i> -acetyl <i>N</i> -glycosyl amides 14a-c , 15a-c , and 16a-c	14
1H , ^{13}C , and NOESY NMR spectra of free <i>N</i> -glycosyl amides 17a-cβ , 18a-cβ , and 19a-cβ	38


Table 1. Chair conformations 4C_1 and 1C_4 and energies obtained for the *xylo* derivates by conformational search employing MMFF94.


Entry	Compound	Conformation 4C_1	Conformation 1C_4
1	$14c\beta$	 18.47 kcal/mol	Not found
2	$15c\beta$	 34.85 kcal/mol	Not found
3	$16c\beta$	 5.82 kcal/mol	Not found
4	$14c\alpha$	 19.71 kcal/mol	 18.25 kcal/mol
5	$15c\alpha$	 36.54 kcal/mol	



			33.99 kcal/mol
6	16ca	Not found	 6.50 kcal/mol


¹H and ¹³C NMR spectra of glycosyl azides **5a-c**.
























^1H and ^{13}C NMR spectra of thioglyco acid 13β and 13α .






















¹H and ¹³C NMR spectra of glycosylamines **2c α** and **2c β** .






-7.26 CDCl₃













^1H , ^{13}C , and NOESY NMR spectra of per-O-acetyl *N*-glycosyl amides **14a-c**, **15a-c**, and **16a-c**.
























































































¹H, ¹³C, and NOESY NMR spectra of free *N*-glycosyl amides **17a-c β** , **18a-c β** , and **19a-c β** .



























-178.4

136.1
134.3
133.1
131.3
131.2
130.7
129.3
128.8
128.4
126.1

~82.4
-79.3
-75.9
-71.9
-71.2

-63.5

-42.3

