

Insights into Structure-Activity Relationships in Efficient Silica-Supported Ni Catalysts for Selective Hydrogenation of Levulinic Acid

Vijaykumar Dosarapu^{a,b}, Siddaramagoud Bandalla^a, Ganesh Babu Bathula^a, Madhu Ravula^a, Satyanarayana Mavurapu^a, Debaprasad Shee^c, Mohan Varkolu^d, Mallesham Baithy^{e,*} and Chandra Shekar Vasam^{a,e,*}

^aDepartment of Pharmaceutical Chemistry, Telangana University, Nizamabad-503322, India

^bDepartment of Chemistry, University College of Science, Satavahana University, Karimnagar-505 002, India

^cChemical Engineering Department, Indian Institute of Technology Hyderabad, Kandi - 502285, Sangareddy, India

^dDepartment of Chemistry, College of Engineering, Koneru Lakshmaiah Education Foundation, R V S Nagar, Aziz Nagar (PO), Moinabad Road, Hyderabad- 500075, India

^eDepartment of Chemistry, School of Science, Gandhi Institute of Technology and Management (GITAM) University, Hyderabad – 502329, Telangana, India

^fDepartment of Chemistry, Guru Ghasidas Vishwavidyalaya (Central University), Bilaspur (CG), India.

*Corresponding Authors:

E-mail: mbaithy@gitam.edu; baithy.m@gmail.com (Dr. Mallesham Baithy)

E-mail: csvasam@telanganauniversity.ac.in (Dr. Chandra Shekar Vasam)




Figure S1. Pyridine FTIR spectra of NiO/HZSM-5 (NiHZSM-5), NiO/SiO₂ (NiSi), NiO/SBA-15 (NISBA-15), NiO/COK-12 (NiCOK-12) catalysts.