

Non-catalytic regioselective synthesis of trans bis-pyrrolo isoxazole cycloadducts

Alhussein Arkan Majhool¹, Assad Abbas Khalaf², Iman Sabeeh Hasan³,
Ranvijay Kumar⁴, Sandeep Kaushal^{5*}, Rahul Badru^{5*}

¹Department of Environmental Health, College of Applied Medical Sciences, University of
Karbala, Karbala, Iraq

²College of Nursing, University of Karbala, Karbala City, Iraq

³Department of Pharmacy, Al-Zahrawi University College, Karbala Iraq

⁴University centre of Research and Development, Chandigarh University, Gharuan, Mohali,
Punjab, India

⁵Department of Chemistry, Sri Guru Granth Sahib World University, Fatehgarh Sahib,
Punjab, India-140406

rahulbadru@gmail.com, kaushalsandeep33@gmail.com

Contents:

¹H-NMR, ¹³C-NMR and FTIR spectra of the synthesized compounds




Figure 1. $^1\text{H-NMR}$ spectrum of compound 3a




Figure 2. $^{13}\text{C-NMR}$ spectrum of compound 3a




Figure 3. FTIR spectrum of compound 3a




Figure 4. ¹H-NMR spectrum of compound 3c




Figure 5. ^{13}C -NMR spectrum of compound 3c




Figure 6. ^1H -NMR spectrum of compound 3g




Figure 7. $^{13}\text{C-NMR}$ spectrum of compound 3g




Figure 8. FTIR spectrum of compound 3g




Figure 9. ^1H -NMR spectrum of compound 3h




Figure 10. ^{13}C -NMR spectrum of compound 3h




Figure 11. ^1H -NMR spectrum of compound 3j




Figure 12. ^{13}C -NMR spectrum of compound 3j




Figure 13. ^1H -NMR spectrum of compound 31




Figure 14. ^{13}C -NMR spectrum of compound 31