


Supporting Information

Artificial viral capsid decorated with DNA aptamer internalizing into lymphoma cells


Kentarou Sakamoto^a, Kohsuke Uchiyama^a, Takashi Iwasaki^b, Hiroshi Inaba^{a,c}, Kazunori Matsuura^{a,c}

- a. Department of Chemistry and Biotechnology, Graduate School of Engineering, Tottori University, Tottori 680-8552, Japan
- b. Department of Bioresources Science, Graduate School of Agricultural Sciences, Tottori University, Tottori 680-8553, Japan
- c. Centre for Research on Green Sustainable Chemistry, Tottori University, Tottori 680-8552, Japan




Supplemental Figure 1. MALDI-TOF MS spectra of (A) β -Annulus-GGGCG, (B) β -Annulus-TE02 and (C) β -Annulus-random DNA.


A




B




C



Supplemental Figure 2. (A) Low-magnification image of 25 μM β -Annulus-TE02, (B) 25 μM β -Annulus and (C) 25 μM TE02 in 10 mM phosphate buffer.



Supplemental Figure 3. Difference spectrum of CD measurement between β -Annulus-TE02 and TE02.



Supplemental Figure 4. Nanoparticle formation of co-assembly of TAMRA- β -Annulus and β -Annulus-TE02. Size distribution (left) and autocorrelation curve (right) obtained from DLS.