## **Supporting information**

Recent advance in optical sensors for microcystin-LR: from recognition elements to signal transduction

Heng Dai<sup>a,1</sup>, Jun Liu<sup>b,1</sup>, Shanlin Wang<sup>c</sup>, Bin Wang<sup>a</sup>, Tianhan Kai<sup>a</sup>, Pian Wu<sup>a\*</sup>, Ping Ding<sup>a\*</sup>

<sup>a</sup> Xiangya School of Public Health, Central South University, Changsha, Hunan, 410013, China

<sup>b</sup> University of South China, Hengyang, Hunan, 421001, China

<sup>c</sup> Guangzhou Customs District, Guangzhou, Guangdong, 510470, China

<sup>1</sup>Heng Dai and Jun Liu contributed equally to this work.

\*Corresponding authors: <u>wupian@csu.edu.cn</u> (P, Wu), <u>pingshui@csu.edu.cn</u> (P, Ding).



Fig. S1. Fundamental steps involving the synthetic imprinting of a polymer to form a MIP.<sup>1</sup>

## **Reference:**

 P. Fernando, M. W. Glasscott, K. Pokrzywinski, B. M. Fernando, G. K. Kosgei and L. C. Moores, *Crit. Rev. Anal. Chem.*, 2022, **52**, 1244-1258.