

Serum IgG galactosylation as a potential biomarker for diagnosis of echinococcosis

Zhuoer Lu,^a Xiaoxiao Feng,^b Bin Fu,^a Xiaojin Mo,^c Ting Zhang,^c Liming Wei,^{*b} Zhonghua Li^{*a,d}
and Haojie Lu^{a,d}

^a Department of Chemistry & Department of Biochemistry and Molecular Biology, School of Basic Medical Sciences, Fudan University, Shanghai 200032, People's Republic of China

^b The Fifth People's Hospital of Shanghai & Institutes of Biomedical Sciences, Fudan University, Shanghai 200032, People's Republic of China

^c National Institute of Parasitic Diseases, Chinese Center for Disease Control and Prevention (Chinese Center for Tropical Diseases Research), Shanghai 200025, People's Republic of China

^d NHC Key Laboratory of Glycoconjugates Research, Fudan University, Shanghai 200032, People's Republic of China

*Corresponding authors.




Figure S1. Five replicate MALDI-TOF-MS spectra of the same sample.




Figure S2. The IgG Gal-ratios of different serum inputs.




Figure S3. The relationship between IgG Gal-ratio and age. The relationship between IgG Gal-ratio and age in control (a), CE (b) and AE (c) patients. Error bars represent SEM of IgG Gal-ratios of the samples at the same age.




Figure S4. Dynamic changes of IgG Gal-ratio with the progression of CE in follow-up patients over three years.