Supramolecular solvent-based ultrasound-assisted extraction of fangchinoline and tetrandrine from the roots of *Stephaniae tetrandrae* 

Yingzhen Wang<sup>a</sup>, Bingyue Cao<sup>a</sup>, Shanshan Shi<sup>a</sup>, Shulin Gao<sup>\*a</sup>, Dandan Ge<sup>b</sup>

<sup>a</sup> School of Chemistry and Chemical Engineering, Kunming University, Kunming

650214, China

<sup>b</sup> School of Chemical Engineering, Yunnan Open University, Yunnan 650223, China

\*Corresponding author. Tel: +86 871 65098043

E-mail address: Shulin Gao, slgao@kmu.edu.cn



Fig. S1 Photos of HFB-hexanol-water system (A) I/L, two-liquid-phase region with the insoluble alkanol in the top and the aqueous phase in the bottom and (B) L/S, liquid-liquid two phase region with aqueous phase in the top and SUPRAS phase in the bottom.



Fig. S2. Representative pictures of the SUPRAS-based UAE procedure (A) Addition of 50 mg of Radix *Stephaniae tetrandrae*, (B) Addition of 600 μL of SUPRAS to the sample, (C) After sonication of the mixture for 50 min, (D) After centrifugation of the mixture.