

Supplementary Information

Acetylated pelargonidin-3-*O*-glucoside alleviates hepatocyte lipid deposition through activating the AMPK-mediated lysosome-autophagy pathway and redox state

Lianghua Xie^{a,b,1}, Xin Hao^{b,1}, Jiahong Xie^b, Jianling Mo^a, Changzheng Yuan^{c*}, Wei Chen^{a,b*}

^a Department of Traditional Chinese Medicine, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou, 310016, China

^b Department of Food Science and Nutrition, College of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou, 310058, China

^c School of Public Health, Zhejiang University School of Medicine, Hangzhou, 310058, China


¹ These authors contributed equally to this work.

*** Corresponding author:**


Prof. Wei Chen, Department of Food Science and Nutrition, College of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou, 310058, China. E-mail: zjuchenwei@zju.edu.cn

Prof. Changzheng Yuan, School of Public Health, Zhejiang University School of Medicine, Hangzhou, 310058, China. E-mail: chy478@zju.edu.cn




1. The raw pictures of Western blot results in Figure 3C






2. The raw pictures of Western blot results in Figure 3E



3. The raw pictures of Western blot results in Figure 4A



4. The raw pictures of Western blot results in Figure 4D



5. The raw pictures of Western blot results in Figure 5A

