

ARTICLE

Table S1. Estimated PAFs (%) that could be potentially contributable to fried food intake from 1997 to 2031 by sex

	Year												
	1997	2000	2004	2006	2009	2011	2013	2016	2018	2022	2025	2029	2031
Male													
CVD	3.2	3.0	4.8	3.6	3.6	3.1	3.9	4.1	4.2	4.5	4.7	4.9	5.0
CHD	2.1	2.0	3.2	2.5	2.4	2.1	2.6	2.8	2.9	3.0	3.1	3.3	3.3
Stroke	13.0	12.8	20.2	15.2	14.8	13.0	16.2	17.2	17.8	19.1	19.9	20.7	21.0
Female													
CVD	2.9	3.4	3.4	3.7	3.1	3.9	4.0	5.1	5.9	7.4	8.5	9.7	10.3
CHD	2.0	2.3	2.3	2.5	2.1	2.6	2.7	3.4	3.9	4.9	5.6	6.5	6.9
Stroke	12.3	14.2	14.3	15.3	12.9	16.1	16.8	21.8	25.2	32.1	36.9	42.4	44.7

Table S2. Estimated PAFs (%) that could be potentially contributable to fried food intake from 1997 to 2031 by age groups

	Year												
	1997	2000	2004	2006	2009	2011	2013	2016	2018	2022	2025	2029	2031
20-29													
CVD	3.3	3.3	4.4	4.0	3.9	4.6	5.1	5.8	6.2	6.8	7.1	7.2	7.2
CHD	2.3	2.2	2.9	2.8	2.6	3.1	3.5	4.0	4.3	4.7	4.9	5.0	5.0
Stroke	13.2	13.9	18.0	16.0	16.0	19.0	20.3	22.8	24.2	26.3	27.1	27.7	27.8
30-39													
CVD	3.3	4.4	3.9	4.1	4.6	4.3	5.5	6.6	7.2	8.6	9.4	10.3	10.6
CHD	2.3	3.0	2.7	2.8	3.1	2.8	3.7	4.4	4.8	5.7	6.2	6.8	7.0
Stroke	13.6	18.3	16.2	16.8	18.9	17.8	23.2	28.1	31.4	37.4	41.2	44.8	46.0
40-49													
CVD	3.3	4.7	4.0	4.3	4.9	4.6	5.5	6.8	7.7	9.3	10.3	11.4	11.8
CHD	2.2	3.1	2.7	2.9	3.3	3.0	3.7	4.6	5.2	6.2	6.9	7.6	7.8
Stroke	13.9	19.7	16.6	17.3	20.2	18.8	23.1	29.0	32.8	39.9	44.5	49.2	51.0
50-59													
CVD	3.1	3.7	4.0	3.8	3.7	4.2	5.1	6.4	7.3	9.1	10.4	11.6	12.1
CHD	2.1	2.5	2.7	2.6	2.5	2.8	3.3	4.3	4.9	6.0	6.8	7.7	8.0
Stroke	12.8	15.5	16.5	15.9	15.2	17.5	21.5	27.6	31.9	40.1	45.4	50.5	52.3
60-69													
CVD	3.3	3.6	4.0	4.1	3.7	4.0	3.7	3.7	3.8	3.8	3.9	3.9	4.0
CHD	2.2	2.4	2.7	2.8	2.5	2.7	2.5	2.5	2.5	2.6	2.6	2.7	2.7
Stroke	13.8	14.8	17.0	17.0	15.4	16.6	15.5	15.6	15.8	16.0	16.2	16.5	16.6
≥70													
CVD	3.0	3.0	4.3	3.5	3.1	3.2	3.7	4.4	4.8	5.6	6.2	6.9	7.2
CHD	2.0	2.0	2.8	2.4	2.1	2.1	2.5	2.9	3.2	3.8	4.2	4.7	4.9
Stroke	12.4	15.5	18.0	14.6	13.2	13.2	15.7	18.4	20.2	23.9	26.6	29.6	30.9

Table S3. Estimated PAFs (%) that could be potentially contributable to fried food intake from 1997 to 2031 by region

	Year												
	1997	2000	2004	2006	2009	2011	2013	2016	2018	2022	2025	2029	2031
Urban													
CVD	4.4	4.5	4.5	5.5	4.5	3.8	4.8	5.5	6.0	6.9	7.5	8.1	8.3
CHD	2.9	3.0	3.0	3.7	3.0	2.6	3.2	3.7	4.1	4.7	5.1	5.5	5.6
Stroke	18.5	18.8	18.7	22.6	18.9	16.1	19.8	23.1	25.3	29.4	31.9	34.3	35.1
Rural													
CVD	2.7	2.8	4.1	3.1	3.0	3.3	3.7	4.3	4.7	5.5	6.1	6.7	7.1
CHD	1.8	1.9	2.7	2.1	2.0	2.2	2.5	2.9	3.1	3.7	4.1	4.5	4.7
Stroke	11.2	11.9	17.3	13.3	12.7	13.9	15.6	18.1	19.9	23.5	26.0	29.0	30.3

ARTICLE




Figure S1. The major types of fried foods included in the CHNS.