

Supporting information

Unveiling the Impact of mpg-C₃N₄@Pa@Ni Nanocomposite in the Reduction of Nitroaromatic Derivatives by Comparative Solvent-free Methods

Fatemeh Eshrati ^a, Hossein Ghafuri*^a, Peyman Hanifehnejad ^a, Haniyeh Dogari ^a

^a Catalysts and Organic Synthesis Research Laboratory, Department of Chemistry, Iran University of Science and Technology, 16846-13114 Tehran, Iran

*E-mail: ghafuri@iust.ac.ir

Contents	Page
1. FT-IR comparison of recovered nanocomposite and origin one's nanocomposite	S2
2. EDS recovered nanocomposite	S3
3. FT-IR spectra of 2-aminobenzophenone	S4
4. ¹ H NMR spectra of 2-aminobenzophenone	S4
5. FT-IR spectra of 4-aminoacetophenone	S5
6. ¹ H NMR spectra of 4-aminoacetophenone	S6
7. FT-IR spectra of 4-bromoaniline	S7
8. ¹ H NMR spectra of 4-bromoaniline	S8




Figure S1. FT-IR comparison of recovered nanocomposite and origin one's nanocomposite.




Figure S2. EDS recovered nanocomposite.




Figure S3. FT-IR spectra of 2-aminobenzophenone.




Figure S4. ^1H NMR spectra of 2-aminobenzophenone. ^1H NMR spectra of 2-aminobenzophenone.




Figure S5. FT-IR spectra of 4-aminoacetophenone.




Figure S6. ¹H NMR spectra of 4-aminoacetophenone.




Figure S7. FT-IR spectra of 4-bromoaniline.




Figure S8. ^1H NMR spectra of 4-bromoaniline.