

Precise control of photoemission and circularly polarized emission achieved through resonant coupling of chiral nanoparticles with Ag particles

Rongze Ma,^a Dedan Mu,^a Yanni Lang,^a Liang Xu,^a Haoxin Wang,^b Songtao Li,^b Yongjin Li,^a Zhiguo Song,^a Dacheng Zhou,^a Yong Yang,^a Yugeng Wen,^a Jin Han,^{*,a} and Jianbei Qiu^{*,a}

^aFaculty of Materials Science and Engineering

Key Laboratory of Advanced Materials of Yunnan Province

Kunming University of Science and Technology

Kunming 650093, P. R. China

Prof. Jin Han, Email: jaakaa@163.com

Prof. Jianbei Qiu, Email: qiu@kust.edu.cn

^bDepartment of Mathematics and Physics, North China Electric Power University,
071003 Baoding, China

Prof. Songtao Li, Email: songtaoli2001@126.com




Figure S1. LCP and RCP UCL from the A-BYE.




Figure S2. Schematic illustration of the energy transfer mechanism.




Figure S3. PL emission spectra of A-BYE samples under 980 nm laser excitation, with full width at half maximum (FWHM) labeled.




Figure S4. PL emission spectra of D-BYE-0.06Ag samples under 980 nm laser excitation, with FWHM labeled.




Figure S5. Dielectric function diagram of D-BYE calculated by first principles.




Figure S6. FDTD solution modeling for D-BYE samples.




Figure S7. FDTD solution modeling for D-BYE-0.06Ag samples.




Figure S8. Electromagnetic field distribution in the X-Y plane obtained from an FDTD simulation of the D-BYE-0.06Ag.




Figure S9. Electromagnetic field distribution in the X-Z plane obtained from an FDTD simulation of the D-BYE-0.06Ag.