

## Supporting Information

### Dual-Band White Light Emission and Temperature-Dependent Luminescence of Sn<sup>2+</sup> in the Metastable Structure of Cs<sub>2</sub>CaCl<sub>4</sub>(H<sub>2</sub>O)<sub>2</sub>


Hu Wang<sup>a</sup>, Yuexiao Pan<sup>a,\*</sup>, Zhenyu Shi<sup>a</sup>, Yelin Cao<sup>a</sup>, Hongzhou Lian<sup>b</sup>, Jun Lin<sup>b,\*</sup>, Qian Miao<sup>a</sup>, Jun Zou<sup>c,d\*</sup>

<sup>a</sup>Key Laboratory of Carbon Materials of Zhejiang Province, College of Chemistry and Materials Engineering, Wenzhou University, Wenzhou 325035, P.R. China.


<sup>b</sup>State Key Laboratory of Rare Earth Resource Utilization, Changchun Institute of Applied Chemistry, Chinese Academy of Sciences, Changchun 130022, P. R. China.

<sup>c</sup>Institute of New Materials & Industrial Technology, Wenzhou University, Wenzhou 325024, China.


<sup>d</sup>School of Science, Shanghai Institute of Technology, Shanghai, 201418, China.




**Fig. S1** Optical images of CCCH:Sn<sup>2+</sup> crystals synthesized from 600 (a), 300 (b), and 100  $\mu\text{L}$  (c) H<sub>3</sub>PO<sub>2</sub> under natural lights..




**Fig. S2** (a) EDS spectrum (inset: SEM) and (b) element proportion of R-CCCH:5%Sn<sup>2+</sup>.




**Fig. S3** Excitation spectra of R-CCCH:xSn<sup>2+</sup> with Sn<sup>2+</sup> ions concentration.




**Fig. S4** Excitation spectra of  $R\text{-CCCH:}x\text{Sn}^{2+}$  varied with  $\text{Sn}^{2+}$  ions concentration.




**Fig. S5** PL spectra of  $R\text{-CCCH:}x\text{Sn}^{2+}$  with different  $\text{Sn}^{2+}$  concentrations.



**Fig. S6** CIE coordinates of  $R\text{-CCCH:}5\%\text{Sn}^{2+}$  crystals excited by different wavelengths.



**Fig. S7** Emission spectra of R-CCCH:Sn<sup>2+</sup> measured at 77 and 298 K under excitation of 284 nm.



**Fig. S8** FWHM of R-CCCH:5%Sn<sup>2+</sup> crystals varied from 77 to 298 K.