

Supporting Information

Secretion analysis of invading process between *Salmonella typhimurium* and HT-29 intestinal epithelial cells on the microfluidic chip

Hong He^{a,c,d}, Haolan Tan^{a,b,d}, Chuang Ge^e, Yi Xu^{a,c,d,*}

^a Key Disciplines Lab of Novel Micro-Nano Devices and System Technology, Key Laboratory of Optoelectronic Technology and Systems, Ministry of Education, Chongqing University, Shapingba, Chongqing 400044, China

^b School of Chemistry and Chemical Engineering, Chongqing University, Shapingba, Chongqing 400044, China

^c School of Optoelectronic Engineering, Chongqing University, Shapingba, Chongqing 400044, China

^d International R & D Center of Micro-nano Systems and New Materials Technology, Chongqing University, Shapingba, Chongqing 400044, China

^e Key Laboratory of Translational Research for Cancer Metastasis and Individualized Treatment, Chongqing University Cancer Hospital, Chongqing 400030, China

* Corresponding author at: Key Disciplines Lab of Novel Micro-Nano Devices and System Technology, Chongqing University, No. 174, St. Shazhengjie, Shapingba District, Chongqing, 400044, China; E-mail address: xuyibbd@cqu.edu.cn (Y. XU)




Fig. S1 The relative cell viability of HT-29 intestinal epithelial cells on the microchip during 4 days cultivation.




Fig. S2 The relative cell viability of HT-29 intestinal epithelial cells within 12 h when co-cultured with *S. typhimurium*