

Supplementary Material

Validation of an RP-HPLC Method for Simultaneously Quantifying Three UV Filters Encapsulated into Nanostructured Lipid Carriers

Andressa C. Schneid^{a†}, Hatylas Azevedo^a, Margarete M. de Araújo^{a†*}

1. Method Development

Table S1. Values of peak area and purity found in DHHB chromatogram.

| Solução | Área | Pureza | % |
|------------------------------------|-----------|--------|--------|
| Standard | 112875027 | 1.00 | 100.00 |
| Standard | 112972317 | 1.00 | 100.09 |
| Standard | 112739741 | 1.00 | 99.88 |
| Ethanol | 0 | 0.00 | 0.00 |
| Acidificated Water:Ethanol (25:75) | 0 | 0.00 | 0.00 |
| Diluent (EtOH+DMF) | 0 | 0.00 | 0.00 |
| Acetonitrile | 0 | 0.00 | 0.00 |
| Blank sample | 0 | 0.00 | 0.00 |
| Blank sample (Filtered) | 0 | 0.00 | 0.00 |

Table S2. Values of peak area and purity found in BEMT chromatogram.

| Solução | Área | Pureza | % |
|------------------------------------|-----------|--------|--------|
| Standard | 499038084 | 1.00 | 100.00 |
| Standard | 499160585 | 1.00 | 100.02 |
| Standard | 498276065 | 1.00 | 99.85 |
| Ethanol | 2306718 | 0.00 | 0.46 |
| Acidificated Water:Ethanol (25:75) | 2418133 | 0.00 | 0.48 |
| Diluent (EtOH+DMF) | 2755972 | 0.00 | 0.55 |
| Acetonitrile | 2684339 | 0.00 | 0.54 |
| Blank sample | 3361542 | 0.00 | 0.67 |
| Blank sample (Filtered) | 2534416 | 0.00 | 0.51 |

Table S3. Values of peak area and purity found in EHT chromatogram.

| Solução | Área | Pureza | % |
|------------------------------------|-----------|--------|--------|
| Standard | 491844734 | 1.0000 | 100.00 |
| Standard | 491961352 | 1.0000 | 100.02 |
| Standard | 489778120 | 1.0000 | 99.58 |
| Ethanol | 2835850 | 1.0000 | 0.58 |
| Acidificated Water:Ethanol (25:75) | 2749282 | 1.0000 | 0.56 |
| Diluent (EtOH+DMF) | 3037644 | 1.0000 | 0.62 |
| Acetonitrile | 3520873 | 1.0000 | 0.72 |
| Blank sample | 3619452 | 1.0000 | 0.74 |
| Blank sample (Filtered) | 3085810 | 1.0000 | 0.63 |




Figure S1. Counter map of light absorption for the standard solution (containing DHHB, BEMT and EHT) detected by DAD in the range of 400 - 800 nm within the time window of 31 minutes.

2. Method Validation

2.1. Linearity

2.1.1. Diethylamino hydroxybenzoyl hexyl benzoate (DHHB)

Table S4. Values concentration and peak area obtained in each level employed in the linearity study for DHHB.

| <i>Sample</i> | <i>Observation</i> | <i>Theoretical Concentration (µg/mL)</i> | <i>Area</i> |
|---------------|--------------------|--|-------------|
| | 1 | 6.0960 | 15035016 |
| Linearity - 1 | 2 | 6.0960 | 15059745 |
| | 3 | 6.0960 | 15058464 |
| | 4 | 12 | 28583242 |
| Linearity - 2 | 5 | 12.0000 | 28635730 |
| | 6 | 12.0000 | 28635730 |
| | 7 | 24.0000 | 56550514 |
| Linearity - 3 | 8 | 24.0000 | 57495099 |
| | 9 | 24.0000 | 56044887 |
| | 10 | 48.0000 | 114510769 |
| Linearity - 4 | 11 | 48.0000 | 113500383 |
| | 12 | 48.0000 | 112871500 |
| | 13 | 60.0000 | 142599948 |
| Linearity - 5 | 14 | 60.0000 | 144725994 |
| | 15 | 60.0000 | 143220791 |
| | 16 | 72.0000 | 170231462 |
| Linearity - 6 | 17 | 72.0000 | 172094837 |
| | 18 | 72.0000 | 172518112 |




Figure S2. Linear fitting for DHHB data obtained in five concentration levels.

Table S5. Data of regression statistics and Cochran Test for DHHB.

| | | |
|--|---------------------------------------|---------------|
| <i>Intercept</i> | | 28922.97741 |
| <i>Inclination</i> | | 2381630.18 |
| <i>Correlation Coefficient:</i> | | 0.999903 |
| <i>Residual Variance</i> | | 7.01098E+11 |
| <i>Standard Error</i> | | 863085.7127 |
| <i>Residual Sum of Squared</i> | | 1.19187E+13 |
| <i>Confidence Interval (95%)</i> | <i>Inferior Limit</i> | -1726171.4253 |
| | <i>Superior Limit</i> | 1726171.4253 |
| | <i>Residual Average</i> | -5.69142E-09 |
| | <i>Standard Deviation of Residues</i> | 837316.1311 |
| | <i>Standardized Inferior Limit</i> | -2.0616 |
| | <i>Standardized Superior Limit</i> | 2.0616 |
| <i>Number</i> | | 18 |
| | <i>N</i> | 16 |
| <i>Cochran Test</i> | | |
| <i>Greater variance of the residuals</i> | | 1.48002E+12 |
| <i>Residuals Variance Sum</i> | | 3.90209E+12 |
| <i>Cochran Statistics</i> | | 0.379 |
| <i>Cochran Critical</i> | | 0.616 |
| <i>Result</i> | | homocedastic |

Table S6. Residue analysis data obtained for the model fitting of DHHB.

| <i>Observation</i> | <i>Y predicted</i> | <i>Residue</i> | <i>Standard Residue</i> | <i>Variance</i> |
|--------------------|--------------------|----------------|-------------------------|-------------------|
| 1 | 14547340.56 | 487675.445 | 0.5824 | |
| 2 | 14547340.56 | 512404.445 | 0.6120 | 193828851.000 |
| 3 | 14547340.56 | 511123.445 | 0.6104 | |
| 4 | 28608485.14 | -25243.138 | -0.0301 | |
| 5 | 28608485.14 | 27244.862 | 0.0325 | 918330048.000 |
| 6 | 28608485.14 | 27244.862 | 0.0325 | |
| 7 | 57188047.3 | -637533.299 | -0.7614 | |
| 8 | 57188047.3 | 307051.701 | 0.3667 | 541835721716.333 |
| 9 | 57188047.3 | -1143160.299 | -1.3653 | |
| 10 | 114347171.6 | 163597.380 | 0.1954 | |
| 11 | 114347171.6 | -846788.620 | -1.0113 | 683929425174.333 |
| 12 | 114347171.6 | -1475671.620 | -1.7624 | |
| 13 | 142926733.8 | -326785.781 | -0.3903 | |
| 14 | 142926733.8 | 1799260.219 | 2.1488 | 1195192282662.330 |
| 15 | 142926733.8 | 294057.219 | 0.3512 | |
| 16 | 171506295.9 | -1274833.941 | -1.5225 | |
| 17 | 171506295.9 | 588541.059 | 0.7029 | 1480016056458.330 |
| 18 | 171506295.9 | 1011816.059 | 1.2084 | |




Figure S3. Plot of residues obtained for each data point from their fitting model for DHHB.



Figure S4. Evaluation of residues dispersion data for DHHB.

Table S7. Outlier evaluation using Z values of each sample. employing Grubb's test for DHHB data.

| <i>Media</i> | 2393155.855 |
|-----------------------------|-------------|
| <i>Standard Deviation</i> | 39363.55528 |
| <i>Z_{critical}</i> | 1.96310 |
| <i>Factor</i> | <i>Z</i> |
| 2466374.016 | 1.8600 |
| 2470430.610 | 1.9631 |
| 2470220.472 | 1.9578 |
| 2381936.833 | 0.2850 |
| 2386310.833 | 0.1739 |
| 2386310.833 | 0.1739 |
| 2356271.417 | 0.9370 |
| 2395629.125 | 0.0628 |
| 2335203.625 | 1.4722 |
| 2385641.021 | 0.1909 |
| 2364591.313 | 0.7257 |
| 2351489.583 | 1.0585 |
| 2376665.800 | 0.4189 |
| 2412099.900 | 0.4813 |
| 2387013.183 | 0.1560 |
| 2364325.861 | 0.7324 |
| 2390206.069 | 0.0749 |
| 2396084.889 | 0.0744 |

2.1.2. Bis-ethylhexyloxyphenol methoxyphenyl triazine (BEMT)

Table S8. Values concentration and peak area obtained in each level employed in the linearity study for BEMT.

| <i>Sample</i> | <i>Observation</i> | <i>Theoretical Concentration ($\mu\text{g/mL}$)</i> | <i>Area</i> |
|---------------|--------------------|--|-------------|
| Linearity - 1 | 1 | 4.0600 | 68170985 |
| | 2 | 4.0600 | 68586504 |
| | 3 | 4.0600 | 68277541 |
| Linearity - 2 | 4 | 8.0000 | 125749366 |
| | 5 | 8.0000 | 127011498 |
| | 6 | 8.0000 | 127011498 |
| Linearity - 3 | 7 | 16.0000 | 251252987 |
| | 8 | 16.0000 | 254160158 |
| | 9 | 16.0000 | 244287700 |
| Linearity - 4 | 10 | 32.0000 | 499897055 |
| | 11 | 32.0000 | 497102551 |
| | 12 | 32.0000 | 492460988 |
| Linearity - 5 | 13 | 40.0000 | 622195704 |
| | 14 | 40.0000 | 632820979 |
| | 15 | 40.0000 | 625475386 |
| Linearity - 6 | 16 | 48.0000 | 751077254 |
| | 17 | 48.0000 | 750181197 |
| | 18 | 48.0000 | 751537867 |




Figure S5. Linear fitting for BEMT data obtained in five concentration levels.

Table S9. Data of regression statistics and Cochran Test for BEMT.

| | | |
|--|---------------------------------------|---------------|
| <i>Intercept</i> | | 2515797.334 |
| <i>Inclination</i> | | 15561197.86 |
| <i>Correlation Coefficient:</i> | | 0.999900 |
| <i>Residual Variance</i> | | 1.382E+13 |
| <i>Standard Error</i> | | 3831942.3011 |
| <i>Residual Sum of Squared</i> | | 2.34941E+14 |
| <i>Confidence Interval (95%)</i> | <i>Inferior Limit</i> | -7663884.6023 |
| | <i>Superior Limit</i> | 7663884.6023 |
| | <i>Residual Average</i> | -4.01504E-08 |
| | <i>Standard Deviation of Residues</i> | 3717530.084 |
| | <i>Standardized Inferior Limit</i> | -2.0616 |
| | <i>Standardized Superior Limit</i> | 2.0616 |
| <i>Number</i> | | 18 |
| | N | 16 |
| <i>Cochran Test</i> | | |
| <i>Greater variance of the residuals</i> | | 2.96018E+13 |
| <i>Residuals Variance Sum</i> | | 7.05021E+13 |
| <i>Cochran Statistics</i> | | 0.420 |
| <i>Cochran Critical</i> | | 0.616 |
| <i>Result</i> | | homocedastic |

Table S10. Residue analysis data obtained for the model fitting of BEMT.

| <i>Observation</i> | <i>Y_{predicted}</i> | <i>Residue</i> | <i>Standard Residue</i> | <i>Variance</i> |
|--------------------|------------------------------|----------------|-------------------------|--------------------|
| 1 | 65694260.65 | 2476724.350 | 0.6662 | 46578059311.000 |
| 2 | 65694260.65 | 2892243.350 | 0.7780 | |
| 3 | 65694260.65 | 2583280.350 | 0.6949 | |
| 4 | 127005380.2 | -1256014.223 | -0.3379 | 530992395141.333 |
| 5 | 127005380.2 | 6117.777 | 0.0016 | |
| 6 | 127005380.2 | 6117.777 | 0.0016 | |
| 7 | 251494963.1 | -241976.113 | -0.0651 | 25738715529562.300 |
| 8 | 251494963.1 | 2665194.887 | 0.7169 | |
| 9 | 251494963.1 | -7207263.113 | -1.9387 | |
| 10 | 500474128.9 | -577073.892 | -0.1552 | 14108075352912.300 |
| 11 | 500474128.9 | -3371577.892 | -0.9069 | |
| 12 | 500474128.9 | -8013140.892 | -2.1555 | |
| 13 | 624963711.8 | -2768007.782 | -0.7446 | 29601753228066.300 |
| 14 | 624963711.8 | 7857267.218 | 2.1136 | |
| 15 | 624963711.8 | 511674.218 | 0.1376 | |
| 16 | 749453294.7 | 1623959.328 | 0.4368 | 475939328653.000 |
| 17 | 749453294.7 | 727902.328 | 0.1958 | |
| 18 | 749453294.7 | 2084572.328 | 0.5607 | |




Figure S6. Plot of residues obtained for each data point from their fitting model for BEMT.




Figure S7. Evaluation of residues dispersion data for BEMT.

Table S11. Outlier evaluation using Z values of each sample. employing Grubb's test for BEMT data.

| | |
|-----------------------------|---------------------|
| Media | 15851127.16 |
| Standard Deviation | 479792.3093 |
| Z_{critical} | 2.17198 |
| Factor | Significance |
| 16790883.005 | 1.9587 |
| 16893227.586 | 2.1720 |
| 16817128.325 | 2.0134 |
| 15718670.750 | 0.2761 |
| 15876437.250 | 0.0528 |
| 15876437.250 | 0.0528 |
| 15703311.688 | 0.3081 |
| 15885009.875 | 0.0706 |
| 15267981.250 | 1.2154 |
| 15621782.969 | 0.4780 |
| 15534454.719 | 0.6600 |
| 15389405.875 | 0.9623 |
| 15554892.600 | 0.6174 |
| 15820524.475 | 0.0638 |
| 15636884.650 | 0.4465 |
| 15647442.792 | 0.4245 |
| 15628774.938 | 0.4634 |
| 15657038.896 | 0.4045 |

2.2.3. Ethylhexyl triazone (EHT)

Table S12. Values concentration and peak area obtained in each level employed in the linearity study for EHT.

| <i>Sample</i> | <i>Observation</i> | <i>Theoretical Concentration (µg/mL)</i> | <i>Area</i> |
|---------------|--------------------|--|-------------|
| Linearity - 1 | 1 | 2.0300 | 70102853 |
| | 2 | 2.0300 | 70993856 |
| | 3 | 2.0300 | 70686250 |
| Linearity - 2 | 4 | 4.0000 | 128881423 |
| | 5 | 4.0000 | 130396028 |
| | 6 | 4.0000 | 130396028 |
| Linearity - 3 | 7 | 8.0000 | 255481127 |
| | 8 | 8.0000 | 259458541 |
| | 9 | 8.0000 | 250291076 |
| Linearity - 4 | 10 | 16.0000 | 508005323 |
| | 11 | 16.0000 | 505068901 |
| | 12 | 16.0000 | 500271395 |
| Linearity - 5 | 13 | 20.0000 | 631220282 |
| | 14 | 20.0000 | 643415347 |
| | 15 | 20.0000 | 633346412 |
| Linearity - 6 | 16 | 24.0000 | 760175508 |
| | 17 | 24.0000 | 759425267 |
| | 18 | 24.0000 | 760956635 |




Figure S8. Linear fitting for EHT data obtained in five concentration levels.

Table S13. Data of regression statistics and Cochran Test for EHT.

| | | |
|--|---------------------------------------|---------------|
| <i>Intercept</i> | | 4591151.375 |
| <i>Inclination</i> | | 31455407.84 |
| <i>Correlation Coefficient:</i> | | 0.999903 |
| <i>Residual Variance</i> | | 1.36366E+13 |
| <i>Standard Error</i> | | 3806433.2729 |
| <i>Residual Sum of Squared</i> | | 2.31823E+14 |
| <i>Confidence Interval (95%)</i> | <i>Inferior Limit</i> | -7612866.5459 |
| | <i>Superior Limit</i> | 7612866.5459 |
| | <i>Residual Average</i> | 9.10627E-09 |
| | <i>Standard Deviation of Residues</i> | 3692782.692 |
| | <i>Standardized Inferior Limit</i> | -2.0616 |
| | <i>Standardized Superior Limit</i> | 2.0616 |
| <i>Number</i> | | 18 |
| | N | 16 |
| <i>Teste de Cochran</i> | | |
| <i>Greater variance of the residuals</i> | | 4.24372E+13 |
| <i>Residuals Variance Sum</i> | | 8.03683E+13 |
| <i>Cochran Statistics</i> | | 0.528 |
| <i>Cochran Critical</i> | | 0.616 |
| <i>Result</i> | | homocedastic |

Table S14. Residue analysis data obtained for the model fitting of EHT.

| Observation | $Y_{predicted}$ | Residue | Standard Residue | Variance |
|--------------------|-----------------------------------|----------------|-------------------------|--------------------|
| 1 | 68445629.29 | 1657223.709 | 0.4488 | 204809976142.334 |
| 2 | 68445629.29 | 2548226.709 | 0.6901 | |
| 3 | 68445629.29 | 2240620.709 | 0.6068 | |
| 4 | 130412782.7 | -1531359.736 | -0.4147 | 764676102008.333 |
| 5 | 130412782.7 | -16754.736 | -0.0045 | |
| 6 | 130412782.7 | -16754.736 | -0.0045 | |
| 7 | 256234414.1 | -753287.097 | -0.2040 | 21133144339370.300 |
| 8 | 256234414.1 | 3224126.903 | 0.8731 | |
| 9 | 256234414.1 | -5943338.097 | -1.6094 | |
| 10 | 507877676.8 | 127646.180 | 0.0346 | 15242046715217.300 |
| 11 | 507877676.8 | -2808775.820 | -0.7606 | |
| 12 | 507877676.8 | -7606281.820 | -2.0598 | |
| 13 | 633699308.2 | -2479026.181 | -0.6713 | 42437248527558.300 |
| 14 | 633699308.2 | 9716038.819 | 2.6311 | |
| 15 | 633699308.2 | -352896.181 | -0.0956 | |
| 16 | 759520939.5 | 654568.458 | 0.1773 | 586351483272.333 |
| 17 | 759520939.5 | -95672.542 | -0.0259 | |
| 18 | 759520939.5 | 1435695.458 | 0.3888 | |




Figure S9. Plot of residues obtained for each data point from their fitting model for EHT.



Figure S10. Evaluation of residues dispersion data for EHT.

Table S15. Outlier evaluation using Z values of each sample. employing Grubb's test for EHT data.

| | |
|-----------------------------|---------------------|
| Media | 31871909.43 |
| Standard Deviation | 435919.8777 |
| Z_{critical} | 1.66796 |
| Factor | Significance |
| 32220355.750 | 0.7993 |
| 32599007.000 | 1.6680 |
| 32599007.000 | 1.6680 |
| 31935140.875 | 0.1451 |
| 32432317.625 | 1.2856 |
| 31286384.500 | 1.3432 |
| 31750332.688 | 0.2789 |
| 31566806.313 | 0.6999 |
| 31266962.188 | 1.3877 |
| 31561014.100 | 0.7132 |

| | |
|--------------|--------|
| 32170767.350 | 0.6856 |
| 31667320.600 | 0.4693 |
| 31673979.500 | 0.4541 |
| 31642719.458 | 0.5258 |
| 31706526.458 | 0.3794 |

2.2.3. LOQ and LOD

Table S16. Signal-to-noise ratio values obtained from the chromatograms for DHHB, BEMT and EHT

| Analyte | Signal/Noise |
|---------|--------------|
| DHHB | 360±18 |
| BEMT | 1014±45 |
| EHT | 1854±69 |

Table S17. LOQ and LOD obtained for DHHB.

| Results | Curve 1 | Curve 2 | Curve 3 | Media | Standard Deviation | LOQ | LOD |
|---------------------|--------------|-------------|-------------|-------------|--------------------|--------|--------|
| Intercept do axis y | 103.967.74 | 230332.6744 | 2694980.771 | | 1460810.669 | 6.0909 | 1.8273 |
| Inclination | 2.370.167.11 | 2396313.455 | 2428603.986 | 2398361.517 | | | |

Table S 19. LOQ and LOD obtained for BEMT.

| Results | Curve 1 | Curve 2 | Curve 3 | Media | Standard Deviation | LOQ | LOD |
|---------------------|---------------|-------------|--------------|-------------|--------------------|--------|--------|
| Intercept do axis y | 1.151.083.02 | 2602903.442 | -11448057.34 | | 7727393.042 | 4.9242 | 1.4773 |
| Inclination da reta | 15.586.228.83 | 15612929.62 | 15879074.78 | 15692744.41 | | | |

Table S 20. LOQ and LOD obtained for EHT.

| Results | Curve 1 | Curve 2 | Curve 3 | Media | Standard Deviation | LOQ | LOD |
|---------------------|---------------|-------------|--------------|-------------|--------------------|--------|--------|
| Intercept do axis y | 3.096.170.70 | 5101720.581 | -7036179.314 | | 6506603.838 | 2.0543 | 0.6163 |
| Inclination da reta | 31.503.927.91 | 31559103.9 | 31956032.87 | 31673021.56 | | | |

2.2.4. Precision (Intra and Inter runs)

Table S21. Precision Intra-Runs DHHB.

| Sample | Concentration (µg/mL) |
|---------------------------|------------------------------|
| Precision Intra 1 | 44.820 |
| Precision Intra 2 | 45.519 |
| Precision Intra 3 | 45.886 |
| Precision Intra 4 | 47.077 |
| Precision Intra 5 | 45.834 |
| Precision Intra 6 | 44.958 |
| Media | 45.682 |
| Standard Deviation | 0.8130 |
| Variance | 0.6609 |
| DPR % | 1.78 |

Table S22. Precision Intra-Runs for BEMT.

| Sample | Concentration (µg/mL) |
|---------------------------|------------------------------|
| Precision Intra 1 | 28.926 |
| Precision Intra 2 | 28.402 |
| Precision Intra 3 | 29.342 |
| Precision Intra 4 | 28.12 |
| Precision Intra 5 | 29.527 |
| Precision Intra 6 | 26.405 |
| Media | 28.453667 |
| Standard Deviation | 1.1380 |
| Variance | 1.2951 |
| DPR % | 4.00 |

Table S23. Precision Intra-Runs for EHT.

| Sample | Concentration (µg/mL) |
|---------------------------|------------------------------|
| Precision Intra 1 | 15.1990 |
| Precision Intra 2 | 15.3260 |
| Precision Intra 3 | 15.6130 |
| Precision Intra 4 | 15.6880 |
| Precision Intra 5 | 15.9490 |
| Precision Intra 6 | 14.7000 |
| Media | 15.412500 |
| Standard Deviation | 0.4392 |
| Variance | 0.1929 |
| DPR % | 2.85 |

Table S24. Precision Inter - Runs for DHHB.

| Sample | Concentration (µg/mL) |
|-------------------|------------------------------|
| Precision Intra 1 | 44.820 |
| Precision Intra 2 | 45.519 |
| Precision Intra 3 | 45.886 |
| Precision Intra 4 | 47.077 |
| Precision Intra 5 | 45.834 |
| Precision Intra 6 | 44.958 |
| Precision Inter 1 | 48.0610 |
| Precision Inter 2 | 46.8510 |
| Precision Inter 3 | 47.3520 |
| Precision Inter 4 | 47.4250 |
| Precision Inter 5 | 47.6410 |
| Precision Inter 6 | 49.5680 |

| | |
|---------------------------|----------------|
| Media | 46.7493 |
| Standard Deviation | 1.3956 |
| Variance (Inter) | 0.8919 |
| DPR % | 2.99 |

Table S25. Precision Inter - Runs for BEMT.

| Sample | Concentration (µg/mL) |
|---------------------------|------------------------------|
| Precision Intra 1 | 15.1990 |
| Precision Intra 2 | 15.3260 |
| Precision Intra 3 | 15.6130 |
| Precision Intra 4 | 15.6880 |
| Precision Intra 5 | 15.9490 |
| Precision Intra 6 | 14.7000 |
| Precision Inter 1 | 16.0020 |
| Precision Inter 2 | 15.5930 |
| Precision Inter 3 | 15.7960 |
| Precision Inter 4 | 15.8860 |
| Precision Inter 5 | 15.9090 |
| Precision Inter 6 | 16.9980 |
| Media | 15.7216 |
| Standard Deviation | 0.5502 |
| Variance (Inter) | 0.2438 |
| DPR % | 3.50 |

Table S26. Precision Inter - Runs for EHT.

| Sample | Concentration (µg/mL) |
|---------------------------|------------------------------|
| Precision Intra 1 | 15.1990 |
| Precision Intra 2 | 15.3260 |
| Precision Intra 3 | 15.6130 |
| Precision Intra 4 | 15.6880 |
| Precision Intra 5 | 15.9490 |
| Precision Intra 6 | 14.7000 |
| Precision Inter 1 | 16.0020 |
| Precision Inter 2 | 15.5930 |
| Precision Inter 3 | 15.7960 |
| Precision Inter 4 | 15.8860 |
| Precision Inter 5 | 15.9090 |
| Precision Inter 6 | 16.9980 |
| Media | 15.7216 |
| Standard Deviation | 0.5502 |
| Variance (Inter) | 0.2438 |
| DPR % | 3.50 |

2.2.5. Accuracy

Table S27. Accuracy for DHHB in NLC.

| Sample | Interval | Standard weight (mg) | Theoretical Concentration (µg/mL) | Experimental concentration (µg/mL) | Recover (%) |
|---------------------------|--------------|----------------------|-----------------------------------|------------------------------------|---------------|
| Accuracy 1A | LOW 80% | 75 | 38.4000 | 38.9090 | 101.33 |
| Accuracy 2A | | 75 | 38.4000 | 39.2010 | 102.09 |
| Accuracy 3A | | 75 | 38.4000 | 37.2620 | 97.04 |
| Accuracy 1B | AVERAGE 100% | 75 | 48.0000 | 48.7980 | 101.66 |
| Accuracy 2B | | 75 | 48.0000 | 47.7340 | 99.45 |
| Accuracy 3B | | 75 | 48.0000 | 48.2290 | 100.48 |
| Accuracy 1C | HIGH 120% | 75 | 57.6000 | 56.9020 | 98.79 |
| Accuracy 2C | | 75 | 57.6000 | 59.0510 | 102.52 |
| Accuracy 3C | | 75 | 57.6000 | 58.0510 | 100.78 |
| Media | | | | | 100.46 |
| Standard Deviation | | | | | 1.76 |
| DPR (%) | | | | | 1.75 |

Table S28. Accuracy for BEMT in NLC

| Sample | Interval | Standard weight (mg) | Theoretical Concentration (µg/mL) | Experimental concentration (µg/mL) | Recover (%) |
|-------------|--------------|----------------------|-----------------------------------|------------------------------------|---------------|
| Accuracy 1A | LOW 80% | 50 | 25.6000 | 26.0420 | 101.73 |
| Accuracy 2A | | 50 | 25.6000 | 27.6750 | 108.11 |
| Accuracy 3A | | 50 | 25.6000 | 24.9200 | 97.34 |
| Accuracy 1B | AVERAGE 100% | 50 | 32.0000 | 33.4860 | 104.64 |
| Accuracy 2B | | 50 | 32.0000 | 32.7490 | 102.34 |
| Accuracy 3B | | 50 | 32.0000 | 32.5600 | 101.75 |
| Accuracy 1C | HIGH 120% | 50 | 38.4000 | 37.9080 | 98.72 |
| Accuracy 2C | | 50 | 38.4000 | 40.2040 | 104.70 |
| Accuracy 3C | | 50 | 38.4000 | 38.9050 | 101.32 |
| | | | | Media | 102.29 |
| | | | | Standard Deviation | 3.24 |
| | | | | DPR (%) | 3.17 |

Table S29. Accuracy for EHT in NLC.

| Sample | Interval | Standard weight (mg) | Theoretical Concentration (µg/mL) | Experimental concentration (µg/mL) | Recover (%) |
|-------------|--------------|----------------------|-----------------------------------|------------------------------------|---------------|
| Accuracy 1A | LOW 80% | 25 | 12.8000 | 13.0250 | 101.76 |
| Accuracy 2A | | 25 | 12.8000 | 13.8440 | 108.16 |
| Accuracy 3A | | 25 | 12.8000 | 12.4930 | 97.60 |
| Accuracy 1B | AVERAGE 100% | 25 | 16.0000 | 16.7150 | 104.47 |
| Accuracy 2B | | 25 | 16.0000 | 16.2260 | 101.41 |
| Accuracy 3B | | 25 | 16.0000 | 16.2300 | 101.44 |
| Accuracy 1C | HIGH 120% | 25 | 19.2000 | 19.0200 | 99.06 |
| Accuracy 2C | | 25 | 19.2000 | 19.9920 | 104.13 |
| Accuracy 3C | | 25 | 19.2000 | 19.3980 | 101.03 |
| | | | | Media | 102.12 |
| | | | | Standard Deviation | 3.12 |
| | | | | DPR (%) | 3.06 |

2.2.6. Robustness

Table S30. Results of Robustness – DHHB.

| Parameters | Analytical method | Changes | k' | Simmetry | Theoretical plates (USP) | Conc. (µg/mL) | Recover |
|--------------------------|--|--|------|----------|--------------------------|---------------|---------|
| Reference values | Alteration | - | 6.16 | 1.18 | 144408 | 49.014 | 102.1 |
| Filter | Filtered | Non Filtered in Filter PVDF 0.45 µm | 6.16 | 1.10 | 143371 | 49.7670 | 103.7 |
| Stability | 0h | 6 days | 6.49 | 1.17 | 139857 | 44.5170 | 92.7 |
| Temperature | 35°C | 34°C | 6.16 | 1.16 | 143952 | 48.961 | 102.0 |
| | | 36°C | 6.14 | 1.13 | 142165 | 48.934 | 101.9 |
| Mobile Phase flow | 0.7mL/min | 0.6mL/min | 6.58 | 1.10 | 153771 | 48.616 | 101.3 |
| | | 0.8 mL/min | 5.83 | 1.21 | 132627 | 48.523 | 101.1 |
| Mobile phase composition | Acidificated Water (H ₃ PO ₄)+EtOH-Scharlau | Acidificated Water (H ₃ PO ₄)+EtOH-Biograde | 6.19 | 1.18 | 146560 | 47.540 | 99.0 |

Table S 31. Results de Robustness – BEMT

| Parameters | Analytical method | Changes | k' | Simmetry | Theoretical plates (USP) | Conc. (µg/mL) | Recover |
|------------|-------------------|---------|------|----------|--------------------------|---------------|---------|
| Reference | Alteration | - | 9.01 | 1.11 | 146045 | 28.6550 | 89.5 |

| | | | | | | | |
|--------------------------|--|--|------|------|--------|---------|-------|
| values | | | | | | | |
| Filter | Filtered | Non Filtered in Filter PVDF 0.45 μm | 9.01 | 1.13 | 146623 | 28.4410 | 88.9 |
| Stability | 0h | 6 days | 9.47 | 1.18 | 147081 | 28.4570 | 88.9 |
| Temperature | 35°C | 34°C | 9.06 | 1.13 | 139750 | 28.624 | 89.5 |
| | | 36°C | 8.95 | 1.15 | 150734 | 28.598 | 89.4 |
| Mobile Phase flow | 0.7mL/min | 0.6mL/min | 7.94 | 1.18 | 254333 | 34.286 | 107.1 |
| | | 0.8 mL/min | 7.31 | 1.22 | 296689 | 34.168 | 106.8 |
| Mobile phase composition | Acidificated Water (H ₃ PO ₄)+EtOH-Scharlau | Acidificated Water (H ₃ PO ₄)+EtOH-Biograde | 7.62 | 1.21 | 266963 | 33.825 | 105.7 |

Table S 32. Results de Robustness – EHT

| Parameters | Analytical method | Changes | k' | Simmetry | Theoretical plates (USP) | Conc. ($\mu\text{g/mL}$) | Recover |
|-----------------|-------------------|--|------|----------|--------------------------|----------------------------|---------|
| Reference value | Alteration | - | 7.58 | 1.16 | 277891 | 16.3690 | 102.3 |
| Filter | Filtered | Non Filtered no Filter PVDF 0.45 μm | 7.58 | 1.18 | 278802 | 16.2360 | 101.5 |
| Stability | 0h | 6 days | 7.98 | 1.19 | 277363 | 24.4490 | 152.8 |

| | | | | | | | |
|--------------------------|--|--|------|------|---------|--------|-------|
| Temperature | 35°C | 34°C | 7.58 | 1.23 | 275378 | 16.342 | 102.1 |
| | | 36°C | 7.56 | 1.23 | 271510 | 16.306 | 101.9 |
| Mobile Phase flow | 0.7mL/min | 0.6mL/min | 9.58 | 1.15 | 1442270 | 14.087 | 88.0 |
| | | 0.8 mL/min | 8.56 | 1.14 | 160150 | 14.057 | 87.9 |
| Mobile phase composition | Acidificated Water (H ₃ PO ₄)+EtOH-Scharlau | Acidificated Water (H ₃ PO ₄)+EtOH-Biograde | 9.20 | 1.12 | 126220 | 13.697 | 85.6 |

2.2.7. System Suitability for Robustness

Table S 33 – System Suitability for Robustness do DHHB

| System Suitability for Robustness | | |
|-----------------------------------|---------|---------|
| Parameter | Minimum | Maximum |
| Capacity factor (k') | 4.31 | 8.01 |
| Assimetry | 0.83 | 1.53 |
| Theoretical plates | 101086 | 187730 |
| Concentration (%) | 46.56 | 51.46 |

Table S34. System Suitability for Robustness do BEMT.

| Parameter | Minimum | Maximum |
|-----------|---------|---------|
|-----------|---------|---------|

| | | |
|----------------------|--------|--------|
| Capacity factor (k') | 6.31 | 11.71 |
| Assimmetry | 0.78 | 1.44 |
| Theoretical plates | 102232 | 189859 |
| Concentration (%) | 27.22 | 30.09 |

Table S35. System Suitability for Robustness do EHT.

| Parameter | Minimum | Maximum |
|----------------------|---------|---------|
| Capacity factor (k') | 5.31 | 9.85 |
| Assimmetry | 0.81 | 1.51 |
| Theoretical plates | 194524 | 361258 |
| Concentration (%) | 15.55 | 17.19 |

3. Response Surface for models of Extraction Content and Encapsulation Efficiency




Figure S11. Response surfaces obtained from prediction models obtained for extraction content of (A) DHHB, (B) BEMT, (C) EHT, and encapsulation efficiency of (D) DHHB, (E) BEMT, (F) EHT.