

Supplementary information

Assessing reactivity of cyanate esters in [2+2+2] cyclootrimerization: Meeting the challenge by augmenting differential scanning calorimetry with kinetic computations

Andrey Galukhin,^{a*} Ilya Nikolaev,^a Alexander Kachmarzhik,^a Roman Nosov,^a Sergey Vyazovkin^{b*}

^a*Alexander Butlerov Institute of Chemistry, Kazan Federal University, 18 Kremlevskaya Street, 420008 Kazan, Russian Federation*

^b*Department of Chemistry, University of Alabama at Birmingham, 901 S. 14th Street, Birmingham, AL 35294, USA*

Corresponding authors:

Andrey Galukhin, Email: and_galuhin@mail.ru

Sergey Vyazovkin, Email: vyazovkin@uab.edu




Figure S1. ¹H NMR spectrum of cyanate ester **1**.




Figure S2. ¹H NMR spectrum of cyanate ester **2**.




Figure S3. ¹H NMR spectrum of cyanate ester **3**.




Figure S4. ^{13}C NMR spectrum of cyanate ester **1**.




Figure S5. ^{13}C NMR spectrum of cyanate ester **2**.




Figure S6. ^{13}C NMR spectrum of cyanate ester **3**.




Figure S7. Chromatogram of cyanate ester **1**.




Figure S8. Chromatogram of cyanate ester **2**.




Figure S9. Chromatogram of cyanate ester **3**.




Figure S10. Raw DSC and baseline curves for cyclotrimerization of cyanate ester **2** at 8 °C min⁻¹.