

Quantum Chemical Prediction and Benchmark Study for Structural Isomers of Terpyridine: Electronic Structure Modulations and Charge Transfer Dynamics

Sehrish Ehsan ^a, Shabbir Muhammad, ^{b*} Shamsa Bibi, ^{a*} Shafiq urRehman, ^a Abdullah G. Al-Sehemi, ^b Abul Kalam, ^b Muhammad Adnan, ^c Muhammad Imran Tousif, ^d

^a Department of Chemistry, University of Agriculture Faisalabad, Faisalabad 38000, Pakistan

^b Department of Chemistry, College of Science, King Khalid University, Abha 61413, P.O. Box 9004, Saudi Arabia

^c Graduate School of Energy Science and Technology, Chungnam National University, Daejeon, 34134, Republic of Korea.

^d Department of Chemistry, Division of Science and Technology, University of Education, 54770 Lahore, Pakistan.

* Corresponding Authors: SM: mshabbir@kku.edu.sa SB: shamsa.shafiq@uaf.edu.pk




Figure S1. Graphical representation of the correlation between the MPP of terpyridine isomers with their excitation energies




Figure S2. Optimized molecular geometries of the studied compounds were obtained at M06-2X/6-311G** calculations.




Figure S3. Graphical representation of UV spectra in THF solvent




Figure S4 Ionization potential tendencies and Electro donating power tendencies under isomerization of terpyridine




Figure S5. ELF (0-1) color-filled maps for TPY-1 to TPY-6 compounds, with color-coded scales: red regions correspond to high electron localization, while blue regions denote delocalized electron density.




Figure S6. LOL (0-0.8) color-filled maps for TPY-1 to TPY-6 compounds, with color-coded scales: red regions correspond to high electron localization, while blue regions denote delocalized electron density.

Table S1. Frontier molecular orbital (FMOs) analysis

Compounds	E_{HOMO}	E_{LUMO}	ΔE	Electronegativity (χ)	Hardness (η)	Chemical potential (μ)
TP-1	-7.67	-0.58	7.09	8.84	4.15	-8.84
TP-2	-7.88	-0.83	7.04	4.34	4.24	-4.34
TP-3	-8.30	-1.07	7.22	4.66	4.32	-4.66
TP-4	-7.90	-0.91	6.99	4.33	4.27	-4.33
TP-5	-7.69	-0.62	7.07	4.16	4.15	-4.16
TP-6	-7.77	-0.82	6.95	4.28	4.19	-4.28
TP-7	-7.91	-0.96	6.96	4.42	4.19	-4.42
TP-8	-7.82	-0.91	6.96	4.27	4.18	-4.27
TP-9	-7.67	-0.63	7.05	4.25	4.15	-4.25
TP-10	-7.81	-0.66	7.14	4.33	4.18	-4.33
TP-11	-8.03	-0.80	7.23	4.40	4.33	-4.40
TP-12	-7.74	-0.73	7.01	4.27	4.23	-4.27

Table S2. Accepting index of terpyridine isomers

Compounds	I	A	ω^-	ω^+
TP-1	8.570	0.270	4.575	8.307
TP-2	8.583	0.100	4.924	8.685
TP-3	8.999	0.340	5.395	9.353
TP-4	8.614	0.064	4.906	8.679
TP-5	8.326	0.007	4.690	8.333
TP-6	8.474	0.091	4.853	8.567
TP-7	8.620	0.226	5.067	8.853
TP-8	8.460	0.081	4.835	8.542
TP-9	8.403	0.097	4.637	8.306
TP-10	8.521	0.145	4.659	8.378
TP-11	8.741	0.067	4.981	8.810

TP-12

8.517

0.040

4.753

8.477

Table S3. Bond length, bond angle and dihedral angle comparison by using M06-2x/6-311G**

Bond length	Computed	Experimental	Difference
C ₃ -C ₅	1.389	1.365	0.024
C ₁ -N ₄₄	1.329	1.330	-0.001
N ₄₄ -C ₉	1.335	1.342	-0.007
C ₉ -C ₁₀	1.492	1.482	0.010
C ₁₀ -N ₄₃	1.333	1.349	-0.016
N ₄₃ -C ₁₆	1.333	1.338	-0.005
C ₁₆ -C ₁₇	1.492	1.490	0.002
C ₁₇ -N ₄₅	1.335	1.330	0.005
N ₄₅ -C ₂₄	1.329	1.338	-0.009
C ₁₃ -C ₂₆	1.482	1.476	0.006
C ₃₁ -C ₄₀	1.420	1.410	0.010
C ₂₆ -C ₂₇	1.422	1.416	0.006
C ₃₄ -C ₃₆	1.415	1.395	0.020
C ₁₈ -C ₂₀	1.386	1.369	0.017
C ₄₀ -C ₄₁	1.416	1.413	0.003
		RMSD	0.019393
Bond angle	Computed	Experimental	Difference
N ₄₄ -C ₉ -C ₁₀	116.6	116.2	0.4
C ₁₀ -N ₄₃ -C ₁₆	118.3	117.5	0.8
N ₄₄ -C ₉ -C ₇	122.5	121.8	0.7
N ₄₃ -C ₁₆ -C ₁₇	116.9	116.3	0.6
C ₁₇ -N ₄₅ -C ₂₄	118.1	117.3	0.8
C ₁₇ -C ₁₈ -C ₂₀	118.6	118.8	-0.2
N ₄₅ -C ₁₇ -C ₁₈	122.6	122.0	-0.6
C ₁₃ -C ₂₅ -C ₂₇	120.0	120.0	0.000
C ₃₈ -C ₄₀ -C ₄₁	121.8	122.2	-0.4
C ₃₁ -C ₄₀ -C ₃₈	118.9	118.7	-0.2

		RMSD		0.530094
Dihedral angle	Computed	Experimental	Difference	
N ₄₄ -C ₉ -C ₁₀ -N ₄₃	178.3	166.7	11.6	
C ₇ -C ₉ -C ₁₀ -C ₁₁	177.9	170.6	7.3	
C ₁₁ -C ₁₃ -C ₂₆ -C ₄₁	143.1	143.5	-0.4	
C ₂₉ -C ₃₁ -C ₄₀ -C ₃₈	179.7	176.1	3.6	
		RMSD		3.93182283

Table S4. Natural bond orbital (NBO) analysis for **TPY-1** to **TPY-6** computed at M06-2X/6-311G** level of theory

Compounds	Donor (i)	Type	Acceptor (j)	Type	E ⁽²⁾ kcal/mol	E(j)-E (i) (a.u.)	F (i;j) (a.u.)
TPY-1	π	C ₁₃ -C ₁₄	π^*	C ₁₆ -N ₄₃	41.72	0.37	0.106
	π	C ₁₃ -C ₁₄	π^*	C ₂₆ -C ₄₁	9.51	0.37	0.055
TPY-2	π	C ₂₆ -C ₃₉	π^*	C ₁₃ -C ₁₄	11.37	0.38	0.06
	σ	C ₁₃ -C ₁₄	σ^*	C ₂₆ -C ₂₇	2.21	1.35	0.049
TPY-3	LP (1)	N ₂₉	π^*	C ₄₀ -C ₄₂	23.72	0.36	0.085
	σ	C ₅₂ -C ₅₄	σ^*	N ₂₉ -C ₅₁	4.97	1.26	0.071
TPY-4	π	C ₁₃ -C ₁₄	π^*	C ₁₆ -N ₂₆	42.33	0.33	0.107
	LP (1)	N ₂₉	π^*	C ₄₀ -C ₄₂	22.31	0.36	0.082
TPY-5	LP (1)	N ₂₉	π^*	C ₄₀ -C ₄₁	23.13	0.36	0.084
	σ	C ₄₀ -C ₄₁	σ^*	C ₄₀ -C ₄₂	5.18	1.41	0.076
TPY-6	LP (1)	N ₇₇	π^*	C ₄₅ -C ₄₇	46.34	0.37	0.122
	π	C ₁₃ -C ₁₄	π^*	C ₁₆ -N ₂₆	42.84	0.33	0.107

Table S5. The centroid distance between two orbitals and HOMO–LUMO orbital overlap integral (S) for **TPY-1** to **TPY-6** computed at M06-2X/6-311G** level of theory

Compounds	Centroid distance (D _{CT})	Orbital overlap integral (S)
TPY-1	3.552	0.657
TPY-2	1.258	0.835
TPY-3	6.296	0.414

TPY-4	6.515	0.398
TPY-5	6.537	0.397
TPY-6	7.491	0.335

Table S6. Benchmark comparison of calculated $\langle\gamma\rangle$ ($\times 10^{-36}$ esu) using different density functionals (B97X-D¹, CAM-B3LYP², LC-BLYP³, M05⁴, M06⁵, and M06-2X⁵) for **TPY-1** to **TPY-6** to assess the influence of range-separation and exchange contribution on the computed results.

Compound	B97X-D/ 6-311G**	CAM-B3LYP/ 6-311G**	M06/ 6-311G**	M05/ 6-311G**	M06-2X/ 6-311G**
TPY-1	59.51	63.21	89.66	86.78	63.83
TPY-2	96.89	105.8	161.0	154.5	108.5
TPY-3	137.0	153.9	266.2	258.2	163.4
TPY-4	161.0	182.4	324.9	314.3	194.7
TPY-5	178.5	201.8	360.8	349.6	219.1
TPY-6	205.9	235.6	410.2	398.5	291.4

Table S7: Literature summary table of $\langle\gamma\rangle$ values for similar terpyridine derivatives

Compounds	NLO values	Reference
Zn(II) complex [Zn(C ₂₇ H ₃₅ N ₃)Cl ₂] with 4,4',4''-tri-tert-butyl-2,2':6',2''-terpyridine ligand	104.19×10^{-36} esu	Hussain S, Muhammad S, Chen X, Akkurt M, Alshehri AM, Din SU, Ullah H, Al-Sehemi AG. Synthesis, Crystal Structure, and Nonlinear Optical Properties of Zn (II) Complex with 4, 4', 4''-Tri-tert-Butyl-2, 2': 6', 2''-Terpyridine: A Dual Exploration. Russian Journal of Inorganic Chemistry. 2020 Mar;65(3):368-77.
[2,6-diphenyl-4-(3,4,5-trimethoxyphenyl) pyridine]	187.53×10^{-36} esu	Chaudhry, A.R., Muhammad, S., Haq, B.U., Kumar, S., Al-Sehemi, A.G., Irfan, A., Laref, A. and Hussain, A., 2019. Exploring the functional properties of Trimethoxy-Phenylpyridine as efficient optical and nonlinear optical material: A quantum chemical approach. <i>Journal of Molecular Structure</i> , 1185, pp.268-275.

References

1. Chai, J.-D. and Head-Gordon, M., *Phys. Chem. Chem. Phys.*, 2008, **10**, 6615.
2. Yanai, T., Tew, D. P. and Handy, N. C., *Chem. Phys. Lett.*, 2004, **393**, 51.
3. Iikura, T., Tsuneda, T., Yanai, T. and Hirao, K., *J. Chem. Phys.*, 2001, **115**, 3540.
4. Zhao, Y., Schultz, N. E. and Truhlar, D. G., *J. Chem. Theory Comput.*, 2005, **1**, 415.
5. Zhao, Y. and Truhlar, D. G., *Theor. Chem. Acc.*, 2008, **120**, 215.