




5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.




5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.




5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.




5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.




5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.



5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.




5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.



5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.





5-ASA\*: 5-ASA group was used as a control for colitis treatment but was excluded from the experimental group as it was not suitable for the purpose of the study to investigate preventive efficacy.