

Supplementary Table S1. Sequences of primers used for RT-qPCR.

Forward (F) and reverse (R) primer sequences for all genes analysed in this study.

Table 1. Oligonucleotide primer sequences used for reverse transcription–quantitative polymerase chain reaction.

Primer name	Primer Sequence (5' to 3')
<i>Nos2</i>	F: CCCTTCAATGGTTGGTACATGG R: ACATTGATCTCCGTGACAGCC
<i>Il1b</i>	F: CTCACAAGCAGAGCACAAGC R: CAGTCCAGCCCATACTTTAGG
<i>Arg1</i>	F: GAACACGGCAGTGGCTTTAAC R: TGCTTAGCTCTGTCTGCTTTGC
<i>Tnf</i>	F: CAGGCGGTGCCTATGTCTC R: CGATCACCCCGAAGTTCAGTAG
<i>Il6</i>	F: CGGAGAGGAGACTTCACAGAG R: CATTTCACGATTTCCCAGA
<i>Plau</i>	F: GCAGCCATCTACCAGAAGAACAAG R: TTGGGAGTTGAATGAAGCAGTGTG
<i>Plat</i>	F: CACTGCCTGCCTCCCTGAC R: GATGCCTCATGCTTGCCGTAG
<i>Bax</i>	F: CGTGGTTGCCCTCTTCTACTTTG R: TCCAGTGTCCAGCCCATGATG
<i>Bcl2</i>	F: TACGAGTGGGATGCTGGAGATG R: TCAGGCTGGAAGGAGAAGATGC
<i>Ldha</i>	F: GGTGGTGGACAGTGCCTACG R: CCCGCCTAAGGTTCTTCATTATGC
<i>Hk2</i>	F: ATCGGCGTGGATGGCTCTG R: GAGGAAGCGGACATCACAATCG
<i>Slc2a1</i>	F: AGAAGAGGGTCGGCAGATGATG R: AGCACCACAGCGATGAGGATG
<i>Pdgfb</i>	F: ACCACTCCATCCGCTCCTTTG R: GTGTGCTCGGGTCATGTTCAAG
<i>β-actin</i>	F: GTGACGTTGACATCCGTAAAGA R: GCCGGACTCATCGTACTCC
<i>Ccl2</i>	F: GCATCTGCCCTAAGGTCTTCAGC

Vegfa

R: AGAAGTGCTTGAGGTGGTTGTGG

F: ACTGGACCCTGGCTTTACTGC

R: TGATCCGCATGATCTGCATGGTG
