

Supporting Information

Exploring metal nanocomposites for asphaltene removal: Role of Cu-BTC, CoMn₂O₄ and K-OMS-2 in adsorption and oxidation of asphaltene

Abhishek Nayak,^a Shanon Viegas,^a Nithya Rajagopal,^a Adrian Marcel Rodrigues,^a Harshini Dasari,^a and Nethaji Sundarabal^{*a}

^a Department of Chemical Engineering, Manipal Institute of Technology, Manipal Academy of Higher Education (MAHE), Manipal, Udipi, Karnataka, India- 576102

Correspondence

Nethaji Sundarabal

Tel: +91 0820 2924316

E-mail: nethaji.s@manipal.edu

List of Figures

Fig.S1 EDX spectrum of A) Cu-BTC, B) CoMn_2O_4 , and C) K-OMS-2.

Fig.S2 EDX spectrum of A) Cu-BTC, B) CoMn_2O_4 , and C) K-OMS-2, after asphaltenes adsorption.

Fig.S3 Effect of adsorbent dosage on asphaltenes adsorption onto A) Cu-BTC (asphaltenes concentration= 250 mg/L, t = 24h, T = 30 °C, 150 rpm), B) CoMn_2O_4 (asphaltenes concentration= 250 mg/L, t = 24h, T = 30 °C, 150 rpm), and C) K-OMS-2 (asphaltenes concentration= 250 mg/L, t = 24h, T = 30 °C, 150 rpm).

Fig.S4 Effect of initial asphaltenes concentration on asphaltenes adsorption onto A) Cu-BTC (adsorbent dosage= 12.5 g/L, t = 24h, T = 30 °C, 150 rpm), B) CoMn_2O_4 (adsorbent dosage= 12.5 g/L, t = 24h, T = 30 °C, 150 rpm), and C) K-OMS-2 (adsorbent dosage= 20 g/L, t = 24h, T = 30 °C, 150 rpm).

Fig.S5 Time dependence studies of asphaltenes adsorption onto A) Cu-BTC (adsorbent dosage= 6.25 g/L, asphaltenes concentration= 500 mg/L, t = 24h, T = 30 °C, 150 rpm), B) CoMn_2O_4 (adsorbent dosage= 6.25 g/L, asphaltenes concentration= 500 mg/L, t = 24h, T = 30 °C, 150 rpm), and C) K-OMS-2 (adsorbent dosage= 6.25 g/L, asphaltenes concentration= 500 mg/L, t = 24h, T = 30 °C, 150 rpm).

Fig.S6 (a) PFO and (b) PSO model plots for the adsorption of asphaltenes onto A) Cu-BTC, B) CoMn_2O_4 , and C) K-OMS-2.

Fig.S7 Intraparticle diffusion model for asphaltenes adsorption onto A) Cu-BTC, B) CoMn_2O_4 , and C) K-OMS-2.

Fig.S8 Boyd model for asphaltenes adsorption onto A) Cu-BTC, B) CoMn_2O_4 , and C) K-OMS-2.

Fig.S9 Langmuir isotherm fitting for asphaltenes adsorption onto A) Cu-BTC (adsorbent dosage= 12.5 g/L, t = 24h, T = 20, 30, 40 °C, 150 rpm), B) CoMn_2O_4 (adsorbent dosage= 12.5 g/L, t = 24h, T = 20, 30, 40 °C, 150 rpm), and C) K-OMS-2 (adsorbent dosage= 20 g/L, t = 24h, T = 20, 30, 40 °C, 150 rpm).

Fig.S10 The van't Hoff plot of asphaltenes adsorption onto A) Cu-BTC, B) CoMn_2O_4 , and C) K-OMS-2.

Fig.S11 The mechanism of asphaltenes adsorption onto Cu-BTC, CoMn_2O_4 , and K-OMS-2.




Fig.S1A




Fig.S1B




Fig.S1C




Fig.S2A




Fig.S2B




Fig.S2C




Fig.S3A




Fig.S3B




Fig.S3C




Fig.S4A




Fig.S4B




Fig.S4C




Fig.S5A




Fig.S5B




Fig.S5C




Fig.S6A




Fig.S6B




Fig.S6C




Fig.S7A




Fig.S7B




Fig.S7C




Fig.S8A




Fig.S8B




Fig.S8C






Fig.S9A






Fig.S9B






Fig.S9C




Fig.S10




Fig.S11