

SUPPLEMENTARY INFORMATION

Self-assembled α/β -dipeptide as dielectrics to improve the behavior of multilayered conducting polymer pseudocapacitors

Dulce A. Quintana-Romero,^{1,2} Adrián Fontana-Escartín,^{1,3,*}

Maria Luisa Gelmi,³ Merve Gul,^{1,3} Maria M. Pérez-Madriral,^{1,3} Raffaella
Bucci,^{2,*} and Carlos Alemán^{1,3,4,*}

¹ *IMEM-BRT Group, Departament d'Enginyeria Química, EEBE, Universitat
Politécnica de Catalunya, C/ Eduard Maristany 10-14, 08019, Barcelona, Spain*

² *Department of Pharmaceutical Sciences, University of Milan, via Venezian 21, Milano
20133, Italy*

³ *Barcelona Research Center in Multiscale Science and Engineering, Campus Diagonal-
Besòs, C/ Eduard Maristany 10-14, 08019, Barcelona, Spain*

⁴ *Institute for Bioengineering of Catalonia (IBEC), The Barcelona Institute of Science
and Technology, Baldiri Reixac 10-12, 08028 Barcelona Spain*





Figure S1. AFM scratch analysis of PEDOT(15s) prepared in acetonitrile. Film thickness was evaluated from height profiles.

Table S1. Conditions used to study the self-assembly of DP1 on PEDOT.

Peptide concentration	Solvent, Cosolvent mixture (v/v %)	Label
0.5 mg/mL	2 % HFIP, 98 % H ₂ O	A
	2 % HFIP, 98 % EtOH	B
	20 % HFIP, 80 % H ₂ O	C
	20 % HFIP, 80 % EtOH	D
2.0 mg/mL	2 % HFIP, 98 % H ₂ O	E
	2 % HFIP, 98 % EtOH	F
	20 % HFIP, 80 % H ₂ O	G
	20 % HFIP, 80 % EtOH	H
	40 % HFIP, 60 % H ₂ O	I
	40 % HFIP, 60 % EtOH	J
	60 % HFIP, 40 % H ₂ O	K
	60 % HFIP, 40 % EtOH	L
	80 % HFIP, 20 % H ₂ O	M
	80% HFIP, 20 % EtOH	N
4.0 mg/mL	40 % HFIP, 60 % H ₂ O	O
	40 % HFIP, 60 % EtOH	P
	60 % HFIP, 40 % H ₂ O	Q
	60 % HFIP, 40 % EtOH	R
8.0 mg/mL	40% HFIP, 60 % H ₂ O	S




Figure S2. Representative optical micrographs of the DP1 assemblies formed using conditions A-D (see Table S1).




Figure S3. Representative optical micrographs of the DP1 assemblies formed using conditions E-N (see Table S1).




Figure S4. Representative optical micrographs of the DP1 assemblies formed using conditions O-R (see Table S1).

Table S2. Conditions used to study the self-assembly of TFA·DP1 on PEDOT.

Peptide concentration	Solvent, Cosolvent mixture (v/v %)	Label
0.5 mg/mL	2 % HFIP, 98 % H ₂ O	A
	40 % HFIP, 60 % H ₂ O	B
	80 % HFIP, 20 % H ₂ O	C
2.0 mg/mL	2 % HFIP, 98 % H ₂ O	D
	2 % HFIP, 98 % EtOH	E
	40 % HFIP, 60 % H ₂ O	F
	40 % HFIP, 60 % EtOH	G
	80 % HFIP, 20 % H ₂ O	H
	80 % HFIP, 20 % EtOH	I
4.0 mg/mL	2 % HFIP, 98 % EtOH	J
8.0 mg/mL	2 % HFIP, 98 % EtOH,	K




Figure S5. Representative optical micrographs of the TFA·DP1 assemblies formed using conditions a-c (see Table S2).




Figure S6. Representative optical micrographs of the TFA·DP1 assemblies formed using conditions d-i (see Table S2).




Figure S7. N 1s high-resolution XPS spectrum for 2PEDOT(3m).




Figure S8. F 1s high-resolution XPS spectrum for (a) 2PEDOT(3m)-TFA·DP1 and (b) 2PEDOT(15s)-TFA·DP1 electrodes.

NMR characterization of TFA·DP1




Figure S9. ¹H NMR spectra of TFA·DP1




Figure S10. ¹³C NMR spectra of TFA·DP1




Figure S11. ^{19}F NMR spectra of TFA·DP1