

## Supplementary Information

### Antibacterial peptidomimetics via fragment display on small-molecule scaffolds

Emma Dyhr<sup>1</sup>, Lucía Cañete de Pinedo<sup>1</sup>, Nicki Frederiksen<sup>1</sup>, Martin Saxtorph Bojer<sup>2</sup>, Hanne Ingmer<sup>2</sup>, Ingvill Pedersen Sæbø<sup>3</sup>, Synnøve Brandt Ræder<sup>3</sup>, Magnar Bjørås<sup>3,4</sup>, Emily Helgesen<sup>3,4</sup>, James Alexander Booth<sup>3,4</sup>, Henrik Franzyk<sup>1</sup>

<sup>1</sup> Department of Drug Design and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, Jagtvej 162, 2100 Copenhagen, Denmark.

<sup>2</sup> Department of Veterinary and Animal Sciences, Section for Bacteria and Viruses, University of Copenhagen, Stigbøjlen 4, 1870 Frederiksberg, Denmark.


<sup>3</sup> Department of Microbiology, Oslo University Hospital and University of Oslo, Rikshospitalet, 0373 Oslo, Norway.

<sup>4</sup> Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology, 7491 Trondheim, Norway.



<b>Contents</b>	<b>Pages</b>
Effect on viability of fibroblast cells	S2-S4
Time-kill experiments	S5-S7
Analytical HPLC and MS data	S7-S26







# Effect on viability of fibroblast cells






Overall comparison of all tested compounds:





Separate experiments:









Time-Kill experiments (compounds **1A**, **1H**, **4E**, **6I**, and **8**)




4E



6I




8




# HPLC and MS

## 1A

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.96$  min. Purity: 99.6%.




**MS:** Calculated for C<sub>108</sub>H<sub>149</sub>F<sub>6</sub>N<sub>21</sub>O<sub>18</sub>S<sub>3</sub>; m/z: [M+2H]<sup>2+</sup> 1119.52392, found 1118.06271




# 1B

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.91$  min. Purity: 97.0%.




**MS:** Calculated for C<sub>111</sub>H<sub>153</sub>F<sub>6</sub>N<sub>21</sub>O<sub>18</sub>S<sub>3</sub>; m/z: [M+2H]<sup>2+</sup> 1140.054739, found 1139.08518




1C

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 10.10$  min. Purity: >99.9%.




**MS:** Calculated for C<sub>114</sub>H<sub>159</sub>F<sub>6</sub>N<sub>21</sub>O<sub>18</sub>S<sub>3</sub>; m/z: [M+2H]<sup>2+</sup> 1161.57087, found 1160.09973




1D

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.70\text{min}$ . Purity: 95.1%.




**MS:** Calculated for C<sub>142</sub>H<sub>198</sub>F<sub>8</sub>N<sub>32</sub>O<sub>20</sub>S<sub>4</sub>; m/z: [M+3H]<sup>3+</sup> 985.14890, found 985.14997




1E

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 10.25$  min. Purity: 99.7%.




**MS:** Calculated for C<sub>74</sub>H<sub>100</sub>F<sub>4</sub>N<sub>14</sub>O<sub>12</sub>S<sub>2</sub>; m/z: [M+2H]<sup>2+</sup> 759.35841, found 758.37461




1H

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.73$  min. Purity: 94.8%.




**MS:** Calculated for C<sub>111</sub>H<sub>156</sub>F<sub>6</sub>N<sub>24</sub>O<sub>21</sub>S<sub>3</sub>; m/z: [M+2H]<sup>2+</sup> 1187.05612, found 1185.57995




## 2D-c1

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.08$  min. Purity: >99.9%.




**MS:** Calculated for C<sub>112</sub>H<sub>150</sub>F<sub>6</sub>N<sub>22</sub>O<sub>18</sub>S<sub>4</sub>; m/z: [M+2H]<sup>2+</sup> 1168.02323, found 1167.03972




## 2D-c2

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.38$  min. Purity: 95.3%.




**MS:** Calculated for C<sub>112</sub>H<sub>150</sub>F<sub>6</sub>N<sub>22</sub>O<sub>18</sub>S<sub>4</sub>; m/z: [M+2H]<sup>2+</sup> 1168.02323, found 1167.03821




### 3D-c1

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.02$  min. Purity: 96.5%.




**MS:** Calculated for C<sub>82</sub>H<sub>110</sub>F<sub>4</sub>N<sub>16</sub>O<sub>14</sub>S<sub>4</sub>; m/z: [M+2H]<sup>2+</sup> 874.36760, found 873.38304




### 3D-c2

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.48$  min. Purity: 96.4%.




**MS:** Calculated for C<sub>82</sub>H<sub>110</sub>F<sub>4</sub>N<sub>16</sub>O<sub>14</sub>S<sub>4</sub>; m/z: [M+2H]<sup>2+</sup> 874.36760, found 873.38251






4F

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.50$  min. Purity: 99.5%.




**MS:** Calculated for C<sub>86</sub>H<sub>122</sub>F<sub>4</sub>N<sub>18</sub>O<sub>13</sub>S<sub>2</sub>; m/z: [M+2H]<sup>2+</sup> 878.44809, found 877.95547




# 4G

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.82$  min. Purity: >99.9%.




**MS:** Calculated for C<sub>86</sub>H<sub>122</sub>F<sub>4</sub>N<sub>18</sub>O<sub>13</sub>S<sub>2</sub>; m/z: [M+2H]<sup>2+</sup> 878.44809, found 877.95491




51

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 8.72$  min. Purity: 97.3%.

**Zoomed Chromatogram ::**




**HR-MS:** Calculated for C<sub>102</sub>H<sub>155</sub>N<sub>21</sub>O<sub>12</sub>; m/z: [M+2H]<sup>2+</sup> 934.61698, found 934.63805




5J

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R$  = 8.70 min. Purity: 90.2%.




**HR-MS:** Calculated for C<sub>102</sub>H<sub>153</sub>N<sub>21</sub>O<sub>12</sub>; m/z: [M+2H]<sup>2+</sup> 933.60933, found 933.60916




5K

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 8.40$  min. Purity: 99.0%.




**HR-MS:** Calculated for C<sub>102</sub>H<sub>156</sub>N<sub>22</sub>O<sub>12</sub>; m/z: [M+2H]<sup>2+</sup> 942.12261, found 942.12265




6I

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.38$  min. Purity: 99.4%.

**Zoomed Chromatogram ::**




**HR-MS:** Calculated for C<sub>102</sub>H<sub>149</sub>F<sub>6</sub>N<sub>21</sub>O<sub>12</sub>; m/z: [M+2H]<sup>2+</sup> 988.58871, found 988.58984




6J

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.37$  min. Purity: 99.6%.




**HR-MS:** Calculated for C<sub>102</sub>H<sub>147</sub>F<sub>6</sub>N<sub>21</sub>O<sub>12</sub>; m/z: [M+2H]<sup>2+</sup> 987.58089, found 987.58163



6K

**Analyt. HPLC:** Buffer A = 5% MeCN + 95% H<sub>2</sub>O + 0.1% TFA. Buffer B = 95% MeCN + 5% H<sub>2</sub>O + 0.1% TFA. Gradient: 0-60% B during 15 min.  $t_R = 9.11$  min. Purity: 99.7%.



**HR-MS:** Calculated for C<sub>102</sub>H<sub>147</sub>F<sub>6</sub>N<sub>21</sub>O<sub>12</sub>; m/z: [M+2H]<sup>2+</sup> 996.09434, found 996.10006

