

Nitrogen cavitation enables rapid and high-yield preparation of functional cell-membrane-derived vesicles

Changting Li^{a,b,†,}, Dehua Luo^{a,†}, Can Peng^a, Qun Luo^a, Wenjing Xu^c, Jincheng Zhou^a,*

Wen Su^c, Wei Wu^d, Yi-Feng Wang^{a,}*

^aGuangzhou Institute of Cancer Research, the Affiliated Cancer Hospital & School of Biomedical Engineering, Guangzhou Medical University, Guangzhou 510260, Guangdong, P.R. China. lichangtingcurie@gmail.com, yifengwang91@gmail.com

^bNewShell VesicleTech of Institut Baulieu, Paris 94270, France..

^cCancer Center, The First Hospital of Jilin University, Changchun 130021, China

^dDepartment of Pathology, Shenzhen University, Shenzhen 518060, China

[†]These authors contributed equally to this work.

Fig. 2e




Fig. 2e




Fig. 2e




Fig. 3c




Fig. 3c




Fig. 3e






Fig. 3e


HEK 293 Δ Nuc
CMDY^{HEK 293} Δ Nuc
HEK 293 Δ Nuc
CMDY^{HEK 293} Δ Nuc



GM130



LaminB1



LAMP1

Fig. 3e






Fig. 5f


CTL
CMDV^{CTL}



LaminB1



LAMP1



VDAC1