

Electronic Supplementary Information


**Hybrid Charge Transfer Complex–Silver Nanoparticles for Portable Colorimetric and Fluorescent Detection of Aflatoxin B1 in Contaminated Peanuts**

Sonam Shakya<sup>a</sup>, Ishaat M. Khan<sup>a\*</sup>, Mohd Anas<sup>a</sup>, Maidul Islam<sup>a,b</sup>, Moamen S. Refat<sup>c</sup>


<sup>a</sup>*Department of Chemistry, Faculty of Science, Aligarh Muslim University, Aligarh-202002, India*

<sup>b</sup>*Research and Development Cell, Lovely Professional University, Phagwara, Punjab-144411, India*


<sup>c</sup>*Department of Chemistry, College of Science, Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia*




**Fig. S1.**  $^1\text{H}$  NMR of (a) ChA, (b) FA, (c) BTy, (d) CTC, and (e) CTAg recorded in  $\text{DMSO-d}_6$ .



**Fig. S2.** TGA-DTA curves for (a) ChA; (b) FA; (c) BTy; (d) CTC; and (e) CTAg.



**Fig. S3.** Job's plot for the CTC-Ag<sup>+</sup> system.



**Fig. S4.** Recyclability of CTAg.

**Table S1.** Elemental percentages (wt%) for CTC and CTAg.

Element	CTC wt%	CTAg wt%
C	69.63	46.23
O	19.18	18.66
Cl	8.27	11.84
N	2.92	1.24
Ag	-	22.03

**Table S2.** Calculation of standard deviation for CTAg

Blank Readings	FL Intensity of CTAg
Reading 1	1226.57
Reading 2	1227.98
Reading 3	1231.01
Reading 4	1232.52
Reading 5	1225.38
<b>Standard Deviation(<math>\sigma</math>)</b>	<b>3.00024499</b>

**Table S3.** Comparison of reported AFB1 Sensors and the present CTag Sensor.

S.No.	Sensing Platform / Material	Detection Method	Detection Limit	Sample Matrix	Reference
1.	Aptamer–MIP with carbon dots & MOF	Fluorescence + Colorimetric	37.0 pg/mL (fluorescence channel) and 13.0 pg/mL (colorimetric channel)	Edible oil	1
2.	Competitive chemiluminescence aptamer	Chemiluminescence	0.09 ng/mL	Wheat	2
3.	Electrochemical aptamer + HCR amplification	Electrochemical	2.84 fg/mL	Corn, medicinal herbs	3
4.	DNAzyme-based colorimetric aptasensor	Colorimetric	3.2 pg/mL	Food samples (general)	4
5.	Smartphone-assisted AuNP aptamer assay	Colorimetric	0.08 ng/g (~80 pg/g)	Beans	5
6.	CTAg (charge-transfer nanomaterial)	Fluorescence + Colorimetric	0.1407 ppb	Peanut	This Work

## References

1. Hu, X., Wang, K., Yang, Y., Ding, B., & Yu, C. (2025). Fluorescence/colorimetric sensor based on aptamers-molecular imprinted polymers synergistic recognition for ultrasensitive and interference-free detection of aflatoxin B1. *Food Chemistry*, 467, 142387.
2. Zhang, Z., Wu, C., & Zhao, Z. (2025). Detection of Aflatoxin B1 in Wheat Based on Nucleic Aptamer Chemiluminescence Sensor. *Sensors*, 25(4), 988.
3. Zhang, H., Ye, S., Huang, L., Fan, S., Mao, W., Hu, Y., ... & Fu, F. (2023). An electrochemical biosensor for the detection of aflatoxin B1 based on the specific aptamer and HCR biological magnification. *Analytical Methods*, 15(1), 99-108.
4. Chen, Y., Zhang, F., Liu, R., Liu, M., Sang, Y., Wang, S., & Wang, X. (2021). A novel colorimetric nano aptasensor for ultrasensitive detection of aflatoxin b1 based on the exonuclease iii-assisted signal amplification approach. *Foods*, 10(11), 2568.

5. Lu, L., Yu, R., & Zhang, L. (2023). AFB1 colorimetric aptamer sensor for the detection of AFB1 in ten different kinds of miscellaneous beans based on gold nanoparticles and smartphone imaging. *Food Chemistry*, 421, 136205.