

Supplementary data

Sustainable and efficient method for sequential extraction of lutein and lipids from deep eutectic solvent pretreated *Chlorella pyrenoidosa*

Beixiao Zhang^{#a*}, Gul Muhammad^{#a,b,c*}, Liya Deng^a, Md Asraful Alam^a, Anqi Zhao^d, Thomas O. Butler^e, Zhenglong Li^{b,c}, Ximing Zhang^{b,c*}, Jingliang Xu^{a*}

^a School of Chemical Engineering, Zhengzhou University, Zhengzhou 450001, China

^b College of Biosystems Engineering and Food Science, Zhejiang University, Hangzhou 310058, China

^c Institute of Zhejiang University-Quzhou, Quzhou 324000, China

^d School of Life Sciences, Zhengzhou University, Zhengzhou 450001, Henan, China.

^e Lgem/Synalgae, Achterweg 65, 1424 PP, De Kwakel, The Netherlands

*Corresponding authors: bxzhang@zzu.edu.cn; gulmuhammad@zju-qz.edu.cn; zhangximing@zju.edu.cn; xujl@zzu.edu.cn

#Both authors contributed equally and will be regarded as 1st co-author




Fig. S1 (a) The cell growth curve of *E. coli* in different media with different supplementary of DES; (b) Staining plot of *E. coli* cells in LB medium under a microscope; (c) Staining plot of *E. coli* cells in LB medium with supplementary of 150 mM DES under a microscope.