

Supporting Information

Synthesis of $\text{Li}_7\text{P}_3\text{S}_{11}$ Solid Electrolyte in Ethyl Propionate Medium for All-Solid-State Li-ion Battery

Tran Anh Tu^{1,2,3} and Nguyen Huu Huy Phuc^{1,2,3*}

¹Faculty of Materials Technology, Ho Chi Minh City University of Technology (HCMUT), 268 Ly Thuong Kiet Str., Dist. 10, Ho Chi Minh City, Vietnam

²National Key Laboratory of Polymer and Composite Materials, Ho Chi Minh City, 268 Ly Thuong Kiet, District 10, Ho Chi Minh City, Vietnam

³Vietnam National University Ho Chi Minh City, Linh Trung Ward, Thu Duc Dist., Ho Chi Minh City, Vietnam

*Corresponding author: nhphuc@hcmut.edu.vn (NHH. Phuc)



Figure S1: Digital photograph of the as-prepared suspension after centrifugation for 5 min at 4000 rpm.




Figure S2: XRD patterns of the precipitate obtained after decanting the yellow supernatant shown in Figure S1 and Li₂S.




Figure S3: Raman spectrum of the precipitate obtained after decanting the yellow supernatant showing in Figure S1; the spectra of EP, Li₂S and P₂S₅ were included for comparison. The inset showed the enlargement of the spectrum of white precipitate.