

Supplementary Information

High-Efficiency Photocatalysis in Ca–Cr Doped BiFeO₃ Nanoparticles via Bandgap tuning

Authors : Anika Rahman Riya ^{a,†}, Abrar Daiyan ^{a,†}, Mehedi Hasan Prince ^{a,b,†}, Troyee Mitra Aishi ^{a,c,†}, Tinni Dey Prome ^a, Md. Abdullah Zubair ^d, Md. Fakhrul Islam ^d, Takian Fakhrul ^{a,*}

Affiliations:

^a Department of Materials and Metallurgical Engineering, Bangladesh University of Engineering and Technology (BUET), Dhaka, 1000, Bangladesh

^b Department of Materials Science & Engineering, Rensselaer Polytechnic Institute (RPI), NY, 12180, USA

^c Department of Chemical Engineering, Northeastern University, MA, 02115, USA

^d Department of Nanomaterials and Ceramic Engineering, Bangladesh University of Engineering and Technology (BUET), Dhaka, 1000, Bangladesh

* Corresponding author. Email: takianf@mme.buet.ac.bd

† Anika Rahman Riya, Abrar Daiyan, Mehedi Hasan Prince, Troyee Mitra Aishi contributed equally to this work

Supplementary information captions:

Figure S1: EDS spectra from multiple points of the BFO sample

Figure S2: EDS spectra from multiple points of the BCFO sample

Figure S3: EDS spectra from multiple points of the BCFOCr1 sample

Figure S4: EDS spectra from multiple points of the BCFOCr3 sample

Figure S5: EDS spectra from multiple points of the BCFOCr5 sample




Figure S1: EDS spectra from multiple points of the BFO sample




Figure S2: EDS spectra from multiple points of the BCFO sample




Figure S3: EDS spectra from multiple points of the BCFOCr1 sample




Figure S4: EDS spectra from multiple points of the BCFOCr3 sample




Figure S5: EDS spectra from multiple points of the BCFOCr5 sample