

Supporting Information for

Locally Ordered Junctions Govern Diffusion in Triglycerides: Insights from Molecular Dynamics Simulations

Yuki Kitamura^a, Yusuke Yasuda^b, Junya Metoki^c, Shogo Tsujino^c, Kazuma Yoshimura^d, Takahide Watanabe^c, and Kazushi Fujimoto^{a,b,e,*}

a) Graduate School of Science and Engineering, Kansai University, 3-3-35 Yamate-cho, Suita, Osaka 564-8680, Japan

b) Organization for Research and Development of Innovative Science and Technology, Kansai University, 3-3-35 Yamate-cho, Suita, Osaka 564-8680, Japan

c) Technical Division, The Nissin OilliO Group, Ltd., 1 Shinmori-cho, Isogo-ku, Yokohama 235-8558, Japan.

d) Household-Use & Wellness Foods Business Strategy, The Nissin OilliO Group, Ltd., 1 Shinmori-cho, Isogo-ku, Yokohama 235-8558, Japan.

e) Department of Chemistry and Materials Engineering, Faculty of Chemistry, Materials and Bioengineering, Kansai University, 3-3-35 Yamate-cho, Suita, Osaka 564-8680, Japan

*Corresponding author. E-mail address: k-fuji@kansai-u.ac.jp

S1 Supporting graph




Figure S1. Potential energy, temperature, and pressure as functions of time for 8:0 (red), 18:1 (blue), and 18:3 (green).