

Table Supplementary 1. Primer sequences used for qRT-PCR

Gene	Forward	Reverse
<i>β-actin</i> (Mouse)	AGAAGATCTGGCACCAC ACC	TACGACCAGAGGCATAC AGG
<i>IL1β</i> (Mouse)	GTGTCTTTCCCGTGGACC TT	AATGGGAACGTCACACA CCA
<i>TNFα</i> (Mouse)	AGCCGATGGGTTGTACC TTG	ATAGCAAATCGGCTGAC GGT
<i>IL6</i> (Mouse)	CTTCTTGGGACTGATGCT GGT	CTCTGTGAAGTCTCCTCT CCG
<i>IL10</i> (Mouse)	TCACTCTTCACCTGCTCC AC	CTATGCTGCCTGCTCTTA CTC
<i>PPARGC</i> <i>1A</i> (Mouse)	GACAGCTTTCTGGGTGG ATT	CGCAGGCTCATTGTTGT ACT
<i>β-actin</i> (Human)	TGACGTGGACATCCGCA AAG	CTGGAAGGTGGACAGCG AGG
<i>PPARGC</i> <i>1A</i> (Human)	TCTGAGTCTGTATGGAG TGACAT	CCAAGTCGTTACATCT AGTTCA