

Supporting Information

Milk exosomes attenuate OVA-induced food allergy by regulating Th1/Th2 balance and remodeling gut microbiota

Siqi Fu^{#1,2,3}, Chao Zhang^{#1,2,3}, Yan Liu^{1,2,3}, Xin Ma^{1,2,3}, Ruofan Xie^{1,2,3}, Hongbing Chen^{1,3,4,5},
and Xin Li^{1,2,3,5*}

¹*State Key Laboratory of Food Science and Resources, Nanchang University, Nanchang 330047, P.R. China.*

²*College of Food Science and Technology, Nanchang University, Nanchang 330047, P.R. China.*

³*Jiangxi Provincial Key Laboratory of Food Allergy, Nanchang University, Nanchang, 330047, P.R. China.*

⁴*Sino German Joint Research Institute, Nanchang University, 235 Nanjing Dong Road, Nanchang, Jiangxi, 330047, China.*

⁵*International Institute of Food Innovation Co., Ltd., Nanchang University, Jiangxi, China*

#These authors contributed equally to this work.

***Corresponding author.**

Xin Li, Professor

School of Food Science and Technology, Nanchang University, Nanchang 330047, China.

Address: No. 235 Nanjing Eastern Road, Nanchang 330047, China

Fax: +86 791 8333708; Tel: +86-791-8830 4447

E-mail address:

Table S1 Primer sequence information

Primer name	Forward primer sequence (5'-3')	Reverse primer sequence (5'-3')
<i>GAPDH</i>	CCTGTTGCTGTAGCCGTATTCA	CCAGGTTGTCTCCTGCGACTT
<i>ZO-1</i>	CGAGGCATCATCCCAAATAAGAA C	TCCAGAAGTCTGCCCGATCAC
<i>Occludin</i>	TTGAAAGTCCACCTCCTTACAGA	CCGGATAAAAAGAGTACGCTGG
<i>MUC2</i>	GGCTGTATTCCCCTCCATCG	CCAGTTGGTAACAATGCCATGT

Fig. S1. Characterization of M-Exo. (A) TEM of M-Exo. (B) The particle size distribution of M-Exo. (C) WB analysis of proteins (CD9, TSG101, CD63 and Calnexin) expression.




Fig. S2 . Demonstration of the original WB strip

