

# Proton-driven Many-Body Interactions and Structural Organization in $\text{He}_n\text{H}^+$ clusters

María Judit Montes de Oca-Estévez<sup>1,†</sup>, Javier Hernández-Rojas<sup>2,†</sup> and Rita  
Prosimi<sup>1\*,†</sup>

<sup>†1</sup> *Institute of Fundamental Physics (IFF-CSIC), CSIC, Serrano 123, 28006 Madrid, Spain*

<sup>‡2</sup> *Departamento de Física and IUdEA, Universidad de La Laguna, 38200 Tenerife, Spain*









E-mail: \*rita@iff.csic.es; Phone: +34-91-5616800Ext.442292





















## SUPPLEMENTARY MATERIAL


















**Table S1:** Calculated relative formation/destruction energies (in  $\text{cm}^{-1}$ ) CCSD(T)/AV6Z level of theory for  $\text{He}_3\text{H}^+$ , taking to account the energy of this system like zero.








Reaction	Relative energy ( $\text{cm}^{-1}$ )
$\text{He}_2\text{H}^+ + \text{He} \rightarrow \text{He}_3\text{H}^+$	355.17
$\text{HeH}^+ + \text{He}_2 \rightarrow \text{He}_3\text{H}^+$	4974.36
$\text{HeH}^+ + 2\text{He} \rightarrow \text{He}_3\text{H}^+$	4981.58
$\text{He}_3 + \text{He}^+ \rightarrow \text{He}_3\text{H}^+$	21429.72
$\text{He}_2 + \text{He} + \text{H}^+ \rightarrow \text{He}_3\text{H}^+$	21437.15
$3\text{He} + \text{H}^+ \rightarrow \text{He}_3\text{H}^+$	21444.38
$\text{He}_3^+ + \text{H} \rightarrow \text{He}_3\text{H}^+$	88594.15
$\text{HeH} + \text{He}_2^+ \rightarrow \text{He}_3\text{H}^+$	90055.51
$\text{He}_2^+ + \text{He} + \text{H} \rightarrow \text{He}_3\text{H}^+$	90060.25
$\text{He}_2 + \text{He}^+ + \text{H} \rightarrow \text{He}_3\text{H}^+$	109987.20
$\text{HeH} + \text{He} + \text{He}^+ \rightarrow \text{He}_3\text{H}^+$	109989.68
$\text{He}^+ + 2\text{He} + \text{H} \rightarrow \text{He}_3\text{H}^+$	109994.42

**Table S2:** Interaction energies (in  $\text{cm}^{-1}$ ) obtained from  $V_{4B}$  potential for  $\text{He}_n\text{H}^+$  complexes, with  $n=3-34$ . The corresponding optimized structures (side and top views) are also depicted.

$[\text{He}_n\text{H}]^+$			This work
$n$	Side view	Top view	$V_{4B}$
3			-21427.5961
4			-21788.9342
5			-22151.5468
6			-22520.6335
7			-22830.5206

8			-23007.2605
9			-23190.1851
10			-23374.7979
11			-23552.5153
12			-23736.0593
13			-23922.4275
14			-24072.7692
15			-24217.6967
16			-24315.9217
17			-24436.2318

18			-24527.3486
19			-24630.3936
20			-24706.6079
21			-24783.0481
22			-24859.2522
23			-24935.2246
24			-25011.5315
25			-25085.1341
26			-25161.5121
27			-25235.1470

28			-25311.7130
29			-25385.0925
30			-25461.7772
31			-25533.5644
32			-25611.8705
33			-25683.1151
34			-25761.3548