

Fe-MOFs-based fluorescent and colorimetric dual-readout nanoprobe for sensitive detection of α -lipomycin

Xianfeng Wang^{1,a}, Genchang Chen^{1,a}, Gongle Shen^{1,a}, Xuefei Chen², Lei Shen³, Bingqing Zhao², Jianhua Su^{4,*}, Dapeng Zhou^{5,6,*}, Sen Yang^{3,*}

¹ Wuxi School of Medicine, Jiangnan University, Wuxi, 214122, PR China

² Department of Clinical Laboratory, Yangpu Hospital, School of Medicine, Tongji University, Shanghai, 200090, China

³ Daqiao Community Healthcare Centre, Yangpu District, Shanghai, 200090, China

⁴ Gerontology, Wuxi No.2 People's Hospital (Jiangnan University Medical Center), PR China

⁵ Life Science and Clinical Medicine Research Center, Affiliated Hospital of Youjiang Medical University for Nationalities, Baise, China

⁶ Department of Burn Plastic and Wound Repair Surgery, Affiliated Hospital of Youjiang Medical University for Nationalities, Baise, China

^a Xianfeng Wang, Genchang Chen and Gongle Shen contributed equally to this work.

* Corresponding authors

E-mail addresses:

suzhou8182@163.com (Jianhua Su)

zhdp1994@163.com (Dapeng Zhou)

ysylq2020@126.com (Sen Yang)




Figure S1 The SEM image (A) and FT-IR (B) of Fe-MOFs after reaction with α -lipomycin.




Figure S2 Effects of response time (A, D) pH (B, E) and temperature (C, F) on the reaction between α -lipomycin and Fe-MOFs nanoprobe in both fluorescent and colorimetric signal.